

BIOSECURITY NEW ZEALAND
IMPORTING COUNTRIES PHYTOSANITARY
REQUIREMENTS
REPUBLIC OF SOUTH AFRICA

Status: Approved

Date: 20 April 1999

Amendment record

Amendment No.	Date:	Nature of amendment:	Approved by:
1.	20 April 1999	Ammendment to sections 2.1, 2.6 and 4.1-4.6 inclusive. Re-issue of EPS.	SCO
2.	26 July1999	Change of address Section 1.1, addition of Section 4.1.2 (Frozen Fruit & Vegetables) and change of headings of Sections 4.6 and 4.6.1 (Herbs.... including Seed for Consumption).	SMN
3.	14 February 2003	Renaming and reformatting of standard. Amendment to Section 2.5 re MPLs.	WJH
4.	1 February 2005	Amendment of MAF contact details Section 1.1 and 1.2. Minor reformatting of document.	WJH
5.	27 March 2007	Amendment of MAF contact details Section 1.1	SW
6.	31 July 2007	Additon of <i>Vaccinium</i> spp. (Blueberry) Commodity specific Requirements (Section 4.1.1); Minor reformatting of the document.	IV
7.	28 July 2008	Interim update: <i>Frankliniella intonsa</i> <i>Tuckerella flabellifera</i> to the quarantine pest list, section 2.2 Additon of <i>Actinidia chinensis</i> (Kiwi fruit) Commodity specific Requirements (Section 4.1.1)	LK
8.	09 October 2008	Correct spelling of <i>Actinidia chinensis</i> (Kiwi fruit) (Section 4.1.1)	LK
9.	20 November 2008	Amendment to remove reference to Wood and Unmanufactured Wood Products; Section 4.7. See Forestry ICPRs. Amendment to MAFBNZ contact details	BHP

Amendment No.	Date:	Nature of amendment:	Approved by:
		Section 1.1	
10.	9 June 2009	<p>Update of contact details for South Africa Plant Health Directorate. NPPOZA June 2009. Section 2.1</p> <p>and</p> <p>Non-commercial imports:</p> <p>Travellers applying for a visa for South Africa must apply well in advance for an import permit from the NPPO South Africa if they wish to bring any agricultural products into South Africa. Visitors must declare any agricultural products and any undeclared products or products without the necessary import permit or phytosanitary certificate will be destroyed. NPPOZA June 2009. Section 2.1</p>	GI

DISCLAIMER

The information in this standard is provided on the following basis. The phytosanitary requirements found in this standard may be used as the basis of export certification. However, requirements may be changed by importing countries at any time at short notice or with no notice to New Zealand. This information is provided strictly on the basis that the Crown, the Ministry of Agriculture & Forestry, its statutory officers, employees, agents and all other persons responsible for or associated with the compilation, writing, editing, approval or publication of the information:

1. disclaim any and all responsibility for any inaccuracy, error, omission, lateness, or any other kind of inadequacy, deficiency or flaw in, or in relation to, the information; and
2. without limiting (1) above, fully exclude any and all liability of any kind on the part of all of them, to any person or entity that chooses to rely on this information

Compliance with this standard is not to be taken as a guarantee that any particular goods will be granted access to any overseas market.

TABLE OF CONTENTS

1	GENERAL INFORMATION.....	5
1.1	FOR ENQUIRES ABOUT THIS STANDARD EMAIL THE EXPORTS GROUP (PLANTS):	5
1.2	PHYTOSANITARY LEGISLATION	5
1.3	DEFINITIONS	5
2.	GENERAL REQUIREMENTS	6
2.1	IMPORT PERMITS	6
2.2	PHYTOSANITARY CERTIFICATES	7
2.3	QUARANTINE PESTS.....	7
2.4	INSPECTION ON ARRIVAL.....	12
2.5	MAXIMUM PEST LIMITS (MPL'S)	12
2.6	PROHIBITIONS.....	12
2.7	PORTS OF ENTRY.....	14
2.8	TRANSIT PROVISIONS.....	14
3.	COMMODITY CLASS REQUIREMENTS	15
4.	COMMODITY SPECIFIC REQUIREMENTS.....	15
4.1	FRUIT AND VEGETABLES	15
4.2	CUT FLOWERS/FOLIAGE.....	20
4.3	MISCELLANEOUS, PACKING MATERIAL AND PLANT (VEGETABLE) FIBRE.....	21
4.4	NURSERY STOCK	22
4.5	SEEDS (GRAIN)/NUTS	65
4.6	HERBS, SPICES, MEDICINAL PLANTS, & PLANTS FOR ANIMAL & HUMAN CONSUMPTION (INCLUDING SEED FOR CONSUMPTION).....	113

1 General Information

1.1 For enquires about this standard email the Plant Exports Group:

plantexports@maf.govt.nz

To help Exports process your email please record in the subject line of your email descriptive keywords which are relevant to your enquiry i.e. ICPR enquiry-Country-specific details.

For urgent enquiries phone or fax the Plant Exports Group
Telephone: 64 4 894 0515
Fax: 64 4 894 0733

1.2 Phytosanitary Legislation

South Africa's phytosanitary requirements are documented in the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992) and is the document on which this export phytosanitary standard is based.

1.3 Definitions

The South Africa Agricultural Pests Act, 1983 (Act No. 36 of 1983) defines the following terms:

Area of production any plot or group of plots that are cultivated as a unit and on which controlled goods are produced;

Certificate the original specimen of a phytosanitary certificate or phytosanitary certificate of re-export that -

- (i) is in the form of the FAO model certificate;
- (ii) has been issued by the recognised authority in the country of production (ie. MAF) or the country of re-export of the controlled goods to which the certificate relates;

Controlled goods (a) any plant, pathogen, insect, exotic animal, growth medium, infectious thing, honey, beeswax or used apiary equipment;

(b) anything determined by the Minister in notice in the Republic of South Africa Government Gazette:

Growth medium any solid or liquid substance in which or on which plants are or can be cultivated, including soil in containers or structures used for cultivation of plants;

Import to introduce controlled goods into the Republic of South Africa or bring about the introduction of such goods into the Republic of South Africa;

Infectious thing	anything except a plant, which may serve as a medium for the importation or spreading of any pathogen, insect or exotic animal;
Insect	any invertebrate member of the animal kingdom irrespective of the stage of development thereof but not such a member - (a) included in the definition of “fish” in section 1 of the Sea Fishery Act, 1988; or (b) that can affect man or an animal only (Definition of “insect” substituted by s. 1 (e) of Act 18 of 1989);
Officer	an officer or employee as defined in section 1 of the Public Service Act, 1984 (Act No. 111 of 1984);
Pathogen	any algae, bacterium, fungus, mycoplasma, rickettsia-like organism, spiroplasma, viroid, or virus, but does not include any such pathogen that can cause a disease in man or an animal only (Definition of “pathogen” substituted by s. 1 (h) of Act 18 of 1989);
Permit	a permit mentioned in section 3 (1) or 6 (3) of the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992);
Plant	any live or dead part of a plant and any derivatives of a plant;
Quarantine area	an area which has been determined as a quarantine area in an order under section 7 of the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992);

2. General Requirements

2.1 Import Permits

Import permits are not required for the commodities specified in section 4.0 of this document unless it is specifically stated. All other commodities require an import permit. Where an import permit is required, phytosanitary information for controlled goods will be specified on the import permit.

Applications for a plant health import permit should be submitted to the Plant Health Import Permit Office at;

Director

Directorate Plant Health

Private Bag X14

Gezina, 0031

OR

Fax: 27 12 3196370

Tel: 27 12 3196102/6396

Email; JeremiahMA@nda.agric.za (as at 9.6.09)

An electronic version of the import permit application form can be accessed from the DoA website at

<http://www.doa.agric.za>

South African NPPO contact:

Director Plant Health

Directorate Plant Health

Department of Agriculture

Private Bag X14

Gezina, 0031

Tel: **27 12 319 6114/6091

Fax: **27 12 319 6580/6101

NPPOZA@nda.agric.za or AliceB@nda.agric.za or Mariannat@nda.agric.za (as at 9.6.09)

Note: Travellers applying for a visa for South Africa must apply well in advance for an import permit from the NPPO South Africa if they wish to bring any agricultural products into South Africa. Visitors must declare any agricultural products and any undeclared products or products without the necessary import permit or phytosanitary certificate will be destroyed. NPPOZA June 2009

2.2 Phytosanitary Certificates

Phytosanitary certificates are required for all commodities unless otherwise specified in section 4.0. Additional declarations and treatments for controlled goods which do not require import permits are specified in section 4.0 (Commodity Class information). For commodities requiring import permits, the phytosanitary requirements are specified on the import permit. Additional declarations and treatments are to be entered in the relevant sections of the phytosanitary certificate.

2.3 Quarantine Pests

The following list of pathogens and pests are prohibited from entering the Republic of South Africa (refer also to the list of prohibited plants and invader plants):-

- | | |
|--|--|
| 1. Abaca mosaic virus | 26. Artichoke curly dwarf virus |
| 2. Abutilon mosaic virus | 27. Artichoke mosaic virus |
| 3. <i>Acalitus essigi</i> | 28. Artichoke mottle crinkle virus |
| 4. <i>Acalitus orthomera</i> | 29. <i>Ascochyta abelmoschi</i> |
| 5. <i>Acarapis woodi</i> | 30. <i>Ascochyta gossypii</i> |
| 6. <i>Aceria biopsida</i> | 31. <i>Ascochyta pisi</i> |
| 7. <i>Aceria breakeyi</i> | 32. <i>Ascochyta rabiei</i> |
| 8. <i>Accria peucedamum</i> | 33. <i>Ascochyta</i> spp. |
| 9. <i>Aceria tulipae</i> | 34. <i>Ascosphaera apis</i> |
| 10. <i>Actinomyces ipomoeae</i> | 35. Asparagus virus |
| 11. <i>Aecidium fragiforme</i> | 36. <i>Aspergillus niger</i> f.sp. <i>floridanus</i> |
| 12. Alfalfa dwarf mosaic virus | 37. <i>Aspidiella hartii</i> |
| 13. <i>Alternaria cichorii</i> | 38. Aster yellows mycoplasma |
| 14. <i>Alternaria padwickii</i> | 39. <i>Bacillus larvae</i> |
| 15. <i>Anarsia lineatella</i> | 40. Bacterial pathogens |
| 16. <i>Anastrepha fraterculus</i> | 41. <i>Balansia oryzae</i> |
| 17. <i>Anastrepha ludens</i> | 42. Banana bunchy top virus |
| 18. <i>Anastrepha mombinpraeoptans</i> | 43. Bean yellow Mosaic virus |
| 19. <i>Anguina</i> spp. | 44. Beet curly top virus |
| 20. <i>Anguina tritici</i> | 45. Black pepper stunt virus |
| 21. Annual ryegrass toxicity syndrome | 46. Blueberry stunt virus |
| 22. <i>Aphelenchoides besseyi</i> | 47. <i>Botrytis tulipae</i> |
| 23. <i>Aphelenchoides fragariae</i> | 48. Broadbean stain virus |
| 24. <i>Aphelenchoides ritzema-bosi</i> | 49. <i>Bursaphelenchus xylophilus</i> |
| 25. Arabis mosaic virus | 50. <i>Cacoecimorpha pronubana</i> |

51. Cadang-cadang viroid
52. *Caecoma sanctae-crucis*
53. *Caecoma torreyae*
54. *Calacarus carinatus*
55. *Campanotus herculeanus*
56. Carnation etched ring virus
57. Carnation necrotic fleck virus
58. Carnation streak virus
59. *Cecidophyopsis ribis*
60. Celery mosaic virus
61. *Cephaleuros parasiticus*
62. *Cephalosporium diospyri*
63. *Cephalosporium maydis*
64. *Cephalosporium sacchari*
65. *Ceratocystis coerulescens*
66. *Ceratocystis fagacearum*
67. *Ceratocystis fimbriata*
68. *Ceratocystis fimbriata* f.sp.
platani
69. *Ceratocystis* spp.
70. *Ceratocystis ulmi*
71. *Ceratostomella fimbriata*
72. *Cercospora angolensis* (Syn.
Phaeoramularia angolensis)
73. *Cercospora coffeicola*
74. *Cercospora corchori*
75. *Cercospora ixorae*
76. *Cercospora mangifera*
77. *Cercospora pini-densiflorae*
78. *Cercospora purpurea*
79. *Cercospora sequioae*
80. *Cercospora sojina*
81. *Cercospora* spp.
82. *Cercospora transversiana*
83. *Ceutospora litchii*
84. *Chaetocnema pulicaria*
85. Cherry leafroll virus
86. Cherry rasp leaf virus
87. Chrysanthemum stunt viroid
88. *Chrysomyxa rhododendri*
89. *Ciboria betulae*
90. Citrus stubborn mycoplasma
91. *Claviceps purpurea*
92. *Claviceps* spp.
93. *Clitocybe tabescens*
94. Closteroviruses
95. Clover phyllody mycoplasma
96. *Cochliobolus miyabeanus* (Syn.
Helminthosporium oryzae)
97. *Colletotrichum cajani*
98. *Colletotrichum capsici*
99. *Colletotrichum coffeanum* (CBD
Strain)
100. *Colletotrichum dematium* f.sp.
truncatum
101. *Colletotrichum fuscum*
102. *Colletotrichum lini*
103. *Colletotrichum panicolae*
104. *Colletotrichum villosum*
105. *Coniella diplodiella*
106. *Corynebacterium flaccumfaciens*
pv. *betae*
107. *Corynebacterium flaccumfaciens*
pv. *flaccumfaciens*
108. *Corynebacterium michiganense*
pv. *michiganense*
109. *Corynebacterium michiganense*
pv. *sepedonicum*
110. *Corynespora asiicola*
111. Cotton leaf curl virus
112. Cranberry false blossom
mycoplasma
113. *Crinipellis palmivora*
114. *Crinipellis perniciosa*
115. *Cronartium coleosporioides*
116. *Cronartium comptoriae*
117. *Cronartium fusiforme*
118. *Cronartium quercuum*
119. *Cronartium strobultinum*
120. *Cryptospora longispora*
121. *Cryptostictis cupressi*
122. Cucumber leafroll virus
123. Cucumber mosaic virus
124. *Cydia molesta*
125. *Dacus cucurbitae*
126. *Dacus dorsalis*
127. *Dacus tryoni*
128. *Dacus zonatus*
129. *Diaporthe phaseolorum* f.sp.
batatatis
130. *Diaporthe phaseolorum* f.sp.
caulivora
131. *Didymella chrysanthemi*
132. *Ditylenchus destructor*
133. *Ditylenchus dipsaci*
134. *Ditylenchus* spp.
135. *Dothistroma pini*
136. *Dotichiza populea*
137. *Drechslera iridis*
138. *Drechslera maydis*
139. *Dysmicoccus brevipes*
140. *Echinodentium taxodii*
141. Elm mosaic virus
142. *Endothia gyrosa*
143. *Endothia parasitica*
144. *Ennomos subsignarius*
145. *Eotetranychus pruni* (*E. pomi*)
146. *Eotetranychus sexmaculatus*
147. *Ephelis pallida*
148. *Eriophyes gastrinichus*
149. *Erwinia amylovora*
150. *Erwinia carotovora* pv. *musae*
151. *Erwinia carotovora* pv. *atroseptica*
152. *Erwinia chrysanthemi*
153. *Erwinia chrysanthemi* pv. *dianthi*
154. *Erwinia nigrifluens*
155. *Erwinia rhapontici*
156. *Erwinia rubrifaciens*
157. *Erwinia stewartii*
158. *Euonymus variegation virus*

159. *Euphorbia* mosaic virus
160. *Exobasidium camelliae*
161. *Exobasidium vexans*
162. *Frankliniella intonsa*
163. *Frankliniella occidentalis*
164. Fungal pathogens
165. *Fusarium oxysporum* f.sp. *cubense*
166. *Fusarium oxysporum* f. sp. *lycopersici* Race III
167. *Fusarium oxysporum* f. sp. *psidii*
168. *Fusarium oxysporum* f.sp. *zingiberi*
169. *Ganoderma lucidum*
170. *Globodera* spp.
171. *Gloeotinia temulenta*
172. *Glomerella cingulata*
173. *Gnomonia platani* (syn *G. veneta*)
174. *Godrona cassandrae*
175. *Goplana dioscorea*
176. *Gremmeniella abeitina*
177. Guar top necrosis virus
178. *Guignardia aesculi*
179. *Guignardia bidwellii*
180. *Gymnosporangium* spp.
181. *Hendersonia agathi*
182. *Heterodera glycines*
183. *Heterodera humuli*
184. *Heterodera* spp.
185. Hop mosaic virus
186. Hop nettle virus complex
187. Hydrangea ringspot virus
188. *Hymenochaeta mongeotii*
189. *Hymenoptera* spp.
190. Iilar group viruses
191. *Kabathina thujae*
192. *Kabatiella nigricans*
193. *Kabatiella zeae*
194. *Keithia thujina*
195. *Koleroga noxia*
196. Leaf scorch
197. *Leptinotarsa decemlineata*
198. *Leptosphaeria maculans* (Syn. *Phoma lingam*)
199. *Leptosphaeria obtusispora*
200. Lethal yellowing mycoplasma
201. Lettuce mosaic virus
202. Lily mosaic virus
203. Lily rosette virus
204. *Liriomyza* spp.
205. *Lophodermium cedri*
206. *Lophodermium* spp.
207. *Lumbricus rubellus*
208. *Lymantria disparina*
209. *Magnaporthe salvinii*
210. *Malpigamoeba mellificae*
211. *Mampava rhodoneura*
212. *Marssonina brannea*
213. *Melampsora hypericorum*
214. *Melampsora pinitorqua*
215. *Meloidogyne* spp.
216. *Microcyclus ulei*
217. *Microsphaeria alni*
218. Mollusca
219. *Monilia roreri*
220. *Monilochaetes infuscans*
221. Mosaic
222. Mosaic virus infection
223. *Mycosphaerella citri*
224. *Mycosphaerella fijiensis*
225. *Mycosphaerella linorum*
226. Myrobolan latent ringspot virus
227. Narcissus yellow stripe virus
228. Necrotic ringspot virus serotypes
229. *Nectria cinnabarina*
230. *Nectria galligena*
231. Nematodes (plant parasitic/plant parasites)
232. *Neovossia indica*
233. *Oidium euonymi-japonici*
234. *Opogona sacchari*
235. *Ovulina azaleae*
236. *Paramyelois transitella*
237. *Pardalaspis cyanescens*
238. Pea seedborne mosaic virus
239. Peanut marginal chlorosis virus
240. Peanut mottle virus (syn. Groundnut mottle virus)
241. Peanut stunt virus
242. Pelargonium leaf curl virus
243. Peperomia ringspot virus
244. *Peronosclerospora maydis*
245. *Peronosclerospora phillipinensis*
246. *Peronosclerospora sacchari*
247. *Peronosclerospora spontanea*
248. *Peridermium cedri*
249. *Peronospora arborescens*
250. *Peronospora documeti*
251. *Peronospora farinosa*
252. *Peronospora hyoschyami*
253. *Peronospora jacksonii*
254. *Peronospora mesembryanthemi*
255. *Peronospora schachtii*
256. *Peronospora tabacina*
257. *Phakopsora cheoana*
258. *Phakopsora jatrophiicola*
259. *Phakopsora pachyrhizi*
260. *Phakopsora zizyphi-vulgaris*
261. *Phoma cisti*
262. *Phoma sabdariffae*
263. *Phoma* spp.
264. *Phoma strasseri*
265. *Phomopsis annonacearum*
266. *Phomopsis heveae*
267. *Phomopsis theae*
268. *Phomopsis vexans*
269. Phormium yellowing mycoplasma
270. *Phyllosticta dracaenae*
271. *Phymatotrichum omnivorum*
272. *Physalospora miyabeana*

273. *Physopella ampelopsidis*
274. Phytophagous mites
275. *Phytophthora capsici*
276. *Phytophthora cryptogea*
277. *Phytophthora fragariae*
278. *Phytophthora infestans*
279. *Phytophthora palmivora*
280. *Phytophthora* spp.
281. Plant parasitic nematodes
282. *Plasmodiophora brassicae*
283. *Plasmopara chrysanthemi-coronarii*
284. *Plasmopara halstedii*
285. *Pleospora papaveracea*
286. *Ploioderma lethale*
287. Plum Pox virus (Sharka)
288. Poinsettia mosaic virus
289. *Polaccia saliciperda*
290. *Polyporus gilvus*
291. *Polyporus sanguineus*
292. *Polyscytalum pustulans*
293. Potato spindle tuber viroid
294. *Pratylenchus brachyurus*
295. *Pratylenchus scribneri*
296. *Prostephanus truncatus*
297. *Pseudococcus citri*
298. *Pseudomonas aleuritides*
299. *Pseudomonas andropogonis*
300. *Pseudomonas aptata*
301. *Pseudomonas caryophylli*
302. *Pseudomonas maublanci*
303. *Pseudomonas phaseolicola*
304. *Pseudomonas saliciperda*
305. *Pseudomonas solanacearum*
306. *Pseudomonas solanacearum* biotypes III and IV
307. *Pseudomonas syringae* pv. *mellea*
308. *Pseudomonas syringae* pv. *pisi*
309. *Pseudomonas syringae* pv. *populae*
310. *Pseudomonas syringae* pv. *savastanoi*
311. *Pseudomonas viridiflava*
312. *Pseudoperonospora humuli*
313. *Psylla pyricola*
314. *Puccinia cari-bistortea*
315. *Puccinia graminis*
316. *Puccinia horiana*
317. *Puccinia psidii*
318. *Puccinia* spp.
319. *Pucciniastrum actinidiae*
320. *Pyrenochaeta phlogina*
321. *Pythium myriotylum*
322. *Radopholus citrophilis*
323. *Radopholus similis*
324. *Ramularia bellulensis*
325. *Ramularia cyclaminicola*
326. *Rhadinaphelenchus cocophilus*
327. *Rhagoletis cerasi*
328. *Rhagoletis pomonella*
329. *Rhizoctonia solani*
330. *Rhyncophorus palmarum*
331. Root wilt
332. Rose Rosette virus
333. Rose wilt virus
334. *Rosselinia bunodes*
335. *Rotylenchus reniformis*
336. *Sanninoidea exitiosa*
337. *Sclerophthora rayssiae* f.sp. *zeae*
338. *Sclerotinia bulborum*
339. *Sclerotinia convulata*
340. *Sclerotinia narcissicola*
341. *Sclerotinia polyblastis*
342. *Sclerotinia ricini*
343. *Sclerotinia trifolium*
344. *Sclerotium tuliparum*
345. *Seiridium cardinale*
346. *Septobasidium aleuritides*
347. *Septoria azaleae*
348. *Septoria gladioli*
349. *Septoria glycines*
350. Shuck die-back disease
351. *Sirex noctilio*
352. Soil-borne viruses
353. *Sorosporium syntherismae*
354. Soybean mild mosaic virus
355. Soybean stunt virus
356. *Sphaceloma manihoticola*
357. *Sphaceloma punicae*
358. *Sphaerotheca mors-uvae*
359. *Sphaerulina taxicola*
360. *Spongospora* spp.
361. *Stenotarsonemus laticeps*
362. Strawberry latent ringspot virus
363. *Streptomyces scabies*
364. Sugarcane Fiji virus
365. Sugarcane grassy shoot virus
366. Sugarcane sereh virus
367. *Synchytrium endobioticum*
368. *Synchytrium piperi*
369. *Tetranychus canadensis*
370. *Tetranychus mcdanielli*
371. *Tetranychus pacificus*
372. *Tetranychus schoenei*
373. *Tetranychus viennensis*
374. *Thyronectria denigrata*
375. *Tilletia barclayana*
376. *Tilletia contraversa*
377. Tobacco leaf curl virus
378. Tobacco rattle virus
379. Tobacco ringspot virus
380. Tobacco streak virus
381. Tomato aspermy virus
382. Tomato black ring virus
383. Tomato ringspot virus
384. *Trioza calacarus*
385. *Trogoderma granarium*
386. *Tropilaelaps clareae*
387. *Tuckerella flabellifera*

- 388 Tulip white streak virus
- 389. Uredinales
- 390. Uredinales (Pine rusts)
- 391. *Uredo phormii*
- 392. *Urocystis agropyri*
- 393. *Urocystis cepulae* (Syn. *U. colchici*)
- 394. *Uromyces cytisi*
- 395. *Uromyces genistae-tinctoriae*
- 396. *Uromyces* spp.
- 397. *Ustilaginoidea virens*
- 398. *Ustilago coicis*
- 399. *Ustilago nuda*
- 400. *Ustilago utriculosa*
- 401. *Ustilago violacea*
- 402. *Varroa jacobsoni*
- 403. *Verticillium albo-atrum*
- 404. *Verticillium dahliae*
- 405. *Verticillium* spp.
- 406. Virus and virus-like diseases affecting honey bees
- 407. Virus and virus diseases
- 408. Virus chlorosis
- 409. Wilt diseases caused by fungi
- 410. *Xanthomonas campestris* pv. *begoniae*
- 411. *Xanthomonas campestris* pv. *campestris*
- 412. *Xanthomonas campestris* pv. *cassavae*
- 413. *Xanthomonas campestris* pv. *cassiae*
- 414. *Xanthomonas campestris* pv. *celebensis*
- 415. *Xanthomonas campestris* pv. *citri*
- 416. *Xanthomonas campestris* pv. *corylina*
- 417. *Xanthomonas campestris* pv. *erythrinae*
- 418. *Xanthomonas campestris* pv. *hyacinthi*
- 419. *Xanthomonas campestris* pv. *khayae*
- 420. *Xanthomonas campestris* pv. *manihotis*
- 421. *Xanthomonas campestris* pv. *oryzae*
- 422. *Xanthomonas campestris* pv. *oryzicola*
- 423. *Xanthomonas campestris* pv. *phormicola*
- 424. *Xanthomonas campestris* pv. *vasculorum*
- 425. *Xanthomonas campestris* pv. *vesicatoria*
- 426. *Xanthomonas fragariae*
- 427. *Xanthomonas gorlincoviae*
- 428. *Xanthomonas* spp.
- 429. *Xanthomonas panici*
- 430. Yam mosaic virus

2.4 Inspection on Arrival

All commodities are subject to inspection on arrival. The level of inspection conducted by South Africa is not known by MAF.

2.5 Maximum Pest Limits (MPL's)

For all commodities exported to South Africa requiring phytosanitary certificates, MPL's are:

Quarantine pests* specified by South Africa	0.5%
	Soil 25g/600unit

*Quarantine pests for South Africa include organisms identified within:

- Section 2.4 of this standard
- Additional declarations
- Phytosanitary import permit

2.6 Prohibitions

Any live plant, vegetative propagative material, or a seed of a kind of plant that has been declared a weed or invader plant is prohibited entry to the Republic of South Africa except by special import permit issued only to a person in charge of a *bona fide* research institution (Agricultural Pets Act 1983 (Act 36 of 1983). Soil (all types) and sand is prohibited from entering the Republic of South Africa.

Declared Weeds include:

Albizia lophantha

Alhagi camelorum

Caesalpinia decapetala

Cannabis sativa

Cereus peruvianus

Cestrum aurantiacum

Cestrum laevigatum

Cestrum parqui

Chromolaena odorata

Cirsium vulgare

Cuscuta campestris

Datura ferox

Datura stramonium

Elodea spp. and other plants of the family

Hydrocharitaceae

Eichhornia spp. and other plants of the family Pontederiaceae

Hakea gibbosa

Hakea sericea

Hakea suaveolens

Harrisia martinii

Lantana camara and any entity which have partly or wholly been derived from the *Lantana camara* complex by means of hybridisation or selection under natural or

artificial conditions.

Lemna spp. and other plants of the family Lemnaceae

Myriophyllum spp. and other plants of the family Haloragidaceae

Opuntia aurantiaca

Opuntia dillenii

Opuntia exaltata

Opuntia ficus-indica excluding all spineless cultivars and selections

Opuntia imbricata

Opuntia lindheimeri

Opuntia rosea

Opuntia spinulifera

Opuntia stricta

Opuntia vulgaris

Orobanche minor

Pereskia aculeata

Pereskia grandifolia

Pereskia stratiotes

Rubus cuneifolius

Salvinia spp. and other plants of the family Salviniaceae

Sesbania punicea

Solanum elaeagnifolium

Solanum mauritianum

Solanum sisymbriifolium
Stipa tenuissima
Stipa trichotoma

Xanthium spinosum
Xanthium strumarium

Declared Invader Plants include:-

Acacia caffra
Acacia cyclops.
Acacia dealbata
Acacia erubescens
Acacia fleckii
Acacia hebeclada
Acacia karroo
Acacia longifolia
Acacia mearnsii
Acacia melanoxylon
Acacia mellifera
Acacia nigrescens
Acacia nilotica
Acacia reficiens subsp. *reficiens*
Acacia robusta subsp. *robusta*
Acacia saligna
Acacia senegal var. *rostrata*
Acacia tenuispina
Acacia tortilis subsp. *heteracantha*
Colophospermum mopane
Combretum apiculatum subsp. *apiculatum*
Commiphora pyracanthoides subsp. *pyracanthoides*
Dichrostachys cinerea subsp. *africana*
Grewia bicolor
Grewia flava
Grewia flavescens
Leptospermum laevigatum
Maytenus senegalensis
Ochna pulchra
Pinus pinaster
Prosopis spp.
Rhigozum trichotomum
Rhus lancea
Tarchonanthus camphoratus
Terminalia sericea

2.7 Ports of Entry

A person importing controlled goods into the Republic of South Africa on the authority of a permit shall do so only through a prescribed port of entry, except where an executive officer has determined some other place.

Prescribed ports include:-

The harbours of: - Cape Town
Durban
East London
Port Elizabeth

The container depots of: - Cape Town
City Deep
Durban
Port Elizabeth

The airports of:- D. F. Malan (Cape Town)
H. F. Verwoerd (Port Elizabeth)
Jan Smuts (Johannesburg)
Lois Botha (Durban)

The border control ports of:- Vioolsdrif
Nakop
Grobler's Bridge
Kopfontein Gate
Schilpad's Gate
Ramatlabama
Beit Bridge
Lebombo
Border Gate (Managa)
Oshoek
Nerston
Mahamba
Golela
Ficksburg Bridge
Maseru Bridge

The Main Post Offices at:- Cape Town
Durban
Johannesburg
East London
Port Elizabeth
Pretoria

2.8 Transit Provisions

The Republic of South Africa has not stipulated transit details for controlled goods.

3. Commodity Class Requirements

Import permits are not required for the commodities specified in section 4.0 unless it is otherwise specified. All other commodities require an import permit. Where an import permit is required phytosanitary information for controlled goods will be specified on the import permit. Phytosanitary certificates are required for all commodities unless otherwise specified. Additional declarations and treatments for controlled goods which do not require import permits are specified in section 4.0 (Commodity Class information). Additional declarations and treatments are to be entered in the relevant sections of the phytosanitary certificate.

4. Commodity Specific Requirements

This section has been split into commodity classes, and further divided into commodity subclasses where necessary. Within each commodity class relevant information on general requirements for import permits (section 2.1), phytosanitary certificates (section 2.2), inspection (section 2.3), prohibitions (section 2.5) and ports of entry (section 2.6) should be referred to with commodity class specific phytosanitary information.

4.1 Fruit and Vegetables

4.1.1 Dried/fresh fruit and vegetables

Does not include seed for consumption or processing.
Phytosanitary import permit required.

Actinidia chinensis

Kiwifruit (fresh fruit)

Conditions:

Phytosanitary import permit required. Phytosanitary certificate required

Additional declaration:

The fruit have been inspected and found free of *Tuckerella flabellifera* (the Peacock mite) and *Frankliniella intonsa* (Eastern Flower thrips).

Vaccinium spp.

Blueberry (fresh fruit)

Conditions:

Phytosanitary import permit required. Phytosanitary certificate required.

Additional declaration:

"The fruit was produced and packed in New Zealand"

and

"New Zealand is free from:

Fungi: *Diaporthe vaccinii*
Godronia cassandrae
Monilinia oxycocci
Monilinia vaccinii-corymbosi

Insects: *Acrobasis vaccinii* [Pyralidae]
Argyrotaenia citrana [Tortricidae]
Argyrotaenia velutinana [Tortricidae]
Ceroplastes floridensis [Coccidae]
Chionaspis salicis [Diaspididae]
Choristoneura rosaceana [Tortricidae]
Conotrachelus nenuphar [Curculionidae]

Eulecanium tiliae [Coccidae]
Grapholita packardi [Tortricidae]
Nemocestes incomptus [Curculionidae]
Pseudococcus maritimus [Pseudococcidae]
Phagoletis mendax [Tephritidae]
Phagoletis tabellaria [Tephritidae]
Rhopobota naevana [Tortricidae]
Sciopithes obscurus [Curculionidae]
Sparganothis sulfureana [Tortricidae]
Spilonota ocellana [Tortricidae]

Mites: *Acalitus vaccinii* [Acari]”

and

“The consignment was inspected, according to the attached Addendum on procedures for inspection, and found free from:

Insects: *Abgrallaspis cyanophylli* [Diaspididae]
Ceroplastes ceriferus [Coccidae]
Ceroplastes sinensis [Coccidae]
Ctenopseustis herana [Tortricidae]
Ctenopseustis obliquana [Tortricidae]
Diaspidiotus ostreaeformis [Diaspididae]
Epiphyas postvittana [Tortricidae]
Otiorhynchus ovatus [Curculionidae]
Otiorhynchus rugosostriatus [Curculionidae]
Otiorhynchus sulcatus [Curculionidae]
Planotortrix excessana [Tortricidae]
Planotortrix octo [Tortricidae]

Mites: *Eotetranychus carpini borealis* [Acari]

ADDENDUM: INSPECTION PROCEDURE

1. Organisms for inspection:

Insects: *Abgrallaspis cyanophylli* [Diaspididae]; *Ceroplastes ceriferus* [Coccidae]; *Ceroplastes sinensis* [Coccidae]; *Ctenopseutis herana* [Tortricidae]; *Ctenopseustis obliquana* [Tortricidae]; *Diaspidiotus ostreaeformis* [Diaspididae]; *Epiphyas postvittana* [Tortricidae]; *Otiorhynchus ovatus* [Curculionidae]; *Otiorhynchus rugosostriatus* [Curculionidae]; *Otiorhynchus sulcatus* [Curculionidae]; *Planotortrix excessana* [Tortricidae]; *Planotortrix octo* [Tortricidae]
Mites: *Eotetranychus carpini borealis* [Acari]

2. Principle of inspection procedure

The principle of inspection according to a specific rate for fruit must be based on a sample of **143** packing units for a consignment of **2000 packing units or less**. The inspection for consignments with **more than 2000 packing units** must be based on **150** packing units. This will provide for a 95% confidence level of detecting packing units with infested fruit if the infestation rate is 2% or higher.

3. Method

3.1 Calculating the sampling interval:

Determine the number of packing units in the consignment intended for export. Divide the number of packing units by 143 or 150 (as determined in point 2). The quotient will be the sampling interval.

3.2 Determining the first packing unit to be inspected:

Randomly select a number from 1 to 13. To this number, add the quotient calculated in point 3.1. This will be the number of first packing unit to be inspected.

3.3 Determining subsequent packing units for inspection:

Add the sampling interval, calculated in point 3.1, to the number of the first packing unit, calculated in point 3.2, to obtain the number of the second packing unit. Determine the number of the third packing unit by adding the number of the second packing unit to the sampling interval. Repeat until the process has accounted for 143 (or 150) packing units.

4. Example for 2 000 packing units: 2 000 packing units ÷ 150 = 13 (13, or the quotient, is the sampling interval); First packing unit to be inspected: select any number from 1 to 13: e.g. 9.; Second packing unit to be inspected: 9 + 13 = 22; Third packing unit to be inspected: 22 + 13 = 35, etc.

5. All fruit from the drawn sample (143 or 150 packing units) shall be inspected and a 5% sample shall be drawn from each packing unit and suspect fruit dissected to determine the status of infestation.

6. Should any of the fruit be found infested with any of the listed quarantine pests, the consignment shall be rejected.

4.1.2 Frozen fruit and vegetables

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Frozen fruit and vegetables require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Actinidia</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , & <i>Dacus dorsalis</i> . <i>Cydia molesta</i> does not occur on kiwifruit in New Zealand. (Please note: This AD to be verified by MAF).
<i>Ananas</i>	AD3:- The area of production is free from <i>Dysmicoccus brevipes</i> AD4: - The consignment is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cydia molesta</i> , & <i>Dacus dorsalis</i> .
<i>Artocarpus</i>	AD2: New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Asimina</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Averrhoa</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .

<i>Bambusa</i>	AD11: - The consignment is free from soil or growth mediums. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Capsicum</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Carica</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> & <i>Dacus tryoni</i> .
<i>Citrus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cercospora angolensis</i> (Syn. <i>Phaeoramularia angolensis</i>), <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> , <i>Mycosphaerella citri</i> & <i>Xanthomonas campestris pv. citri</i> .
<i>Citrus</i> (for Jewish festivals)	No additional declarations required. T8 = the consignment was treated by immersion in sodium hypochlorite. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Citrus</i> (Candied peel)	AD2: - New Zealand is free from <i>Cercospora angolensis</i> (Syn. <i>Phaeoramularia angolensis</i>), and <i>Xanthomonas campestris pv. citri</i> .
<i>Cocos</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Colchicum</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
Cucurbitaceae	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> & <i>Dacus tryoni</i> .
<i>Curcuma</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Dioscorea</i>	AD11: - The consignment is free from soil or growth mediums. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Diospyros</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Fragaria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .
<i>Litchi</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .

<i>Lycopersicon</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , & <i>Pardalaspis cyanescens</i> .
<i>Malus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , & <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> , <i>Eotetranychus sexmaculatus</i> & <i>Rhagoletis pomonella</i> .
<i>Manilkara</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Myricaria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Olea</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Passiflora</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Persea</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Phaseolus</i>	Phytosanitary certificate required.
<i>Phoenix</i> (with seed)	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Phoenix</i> (stoned)	Phytosanitary certificate required.
<i>Physalis</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Pisum</i>	Phytosanitary certificate required.
<i>Pouteria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Psidium</i>	AD2: - New Zealand is free from <i>Aceria biopsida</i> , <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> & <i>Eotetranychus sexmaculatus</i> . AD3:- The area of production is free from <i>Puccinia psidii</i> . AD4: - The consignment is free from <i>Cydia molesta</i> .
<i>Punica</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .

<i>Pyrus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
<i>Ribes</i>	AD2: - New Zealand is free from <i>Aceria breakeyi</i> , <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cecidophyopsis ribis</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .
Rosaceae (Genera & sp. not mentioned elsewhere)	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
<i>Rubus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Mycosphaerella citri</i> . AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
Rutaceae	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Mycosphaerella citri</i> & <i>Xanthomonas campestris</i> pv. <i>citri</i> .
<i>Solanum</i> (excluding <i>S.</i> <i>tuberosum</i>)	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Zea</i>	Phytosanitary certificate required.
<i>Zizyphus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Rhagoletis cerasi</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> .

4.2 Cut Flowers/Foliage

4.2.1 Dried/fresh cut flowers/foliage

Phytosanitary import permit required.

4.3 Miscellaneous, Packing Material and Plant (Vegetable) Fibre

4.3.1 Miscellaneous and packing material

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Miscellaneous and packing material require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7).

Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

Compost, compost inoculum, vermi compost	AD:- Either "Laboratory examination of representative samples has shown the consignment to be free from plant pathogenic - bacteria, fungi, nematodes and soil." or "The consignment has been sterilised."
Bamboo/Rattan	No additional declarations required.
Bark (of <i>Barcena</i> spp. only)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
Cork (unmanufactured)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
Cork (manufactured)	No additional declarations required.
Growth media & Orchid rooting media	(excluding bark of coniferous plants).AD:- Either "Laboratory examination of representative samples has shown the consignment to be free from plant pathogenic - bacteria, fungi, nematodes and soil." or "The consignment has been sterilised."
Resins/Vegetable gums	No additional declarations required.
Sawdust	Import permit required.
Tobacco (leaf, stalks etc.)	Import permit required.

4.3.2 Plant (Vegetable) Fibre

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Plant (Vegetable) Fibre require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

Bombacaceae	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. Must contain less than 1 seed per 500g fibre.
Cocos	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).
<i>Corchoris</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).

<i>Gossypium</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. Must contain less than 1 seed per 2kg fibre.
<i>Hibiscus</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
<i>Leopoldiana piassuba</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).
<i>Linum</i>	No additional declarations required.
<i>Sorghum</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.

4.4 Nursery Stock

4.4.1 Budwood/Cuttings

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Bud wood/Cuttings require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Abelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acalypha</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acanthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Achillea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acokanthera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Actinidia</i>	AD4:- The consignment is free from <i>Eotetranychus sexmaculatus</i> , <i>Pseudomonas viridiflava</i> and <i>Pucciniastrum actinidiae</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Actinodium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Adenanthos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Adenia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.

<i>Adenium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Agave</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , and <i>Steneotarsonemus laticeps</i> . T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aglaonema</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ajuga</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Alberta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aleurites</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas aleuritides</i> , <i>Septobasidium aleuritides</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aloe</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Polyporus sanguineus</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Amaranthaceae	AD4:- The consignment is free from Beet curly top virus, Cucumber mosaic virus, Tobacco ringspot virus, Tomato black ring virus and Tomato ringspot virus.
<i>Amphicarpaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Androsace</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Angelica</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Anigozanthos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Anthurium</i>	AD4:- The consignment is free from <i>Radopholus citrophilis</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Antigonon</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aphelandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aquilegia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, & Cucumber mosaic virus did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.

<i>Arabis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria maculans</i> (Syn. <i>Phoma lingam</i>) & <i>Xanthomonas campestris</i> pv. <i>campestris</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aralia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Arbutus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ardisia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Arisaema</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aristolochia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Armeria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Armoracia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Celery mosaic virus, <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Artemisia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Plasmopara chrysanthemi-coronarii</i> , & <i>Plasmopara halstedii</i> , did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Artocarpus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Koleroga noxia</i> & <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Asterosperma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Atropa</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora hyoschymi</i> , & Tobacco rattle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Azadirachta</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ganoderma lucidum</i> & <i>Polyporus gilvus</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Azalea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Baeckea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Banksia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Barcena</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bauera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bauhinia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Beaucarnea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Beaufortia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Belaperone</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bellis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Bischofia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bougainvillea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bouvardia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Brachychiton</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Brighamia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Buddleia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calathea</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> .
<i>Callistemon</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Callitris</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Cercospora sequioae</i> , <i>Cryptostictis cupressi</i> , <i>Gymnosporangium</i> spp., <i>Kabathina thujae</i> , & <i>Seiridium cardinale</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Calothamnus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Calpurnia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calycanthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calytrix</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Camellia</i>	AD2:- The area of production is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , & <i>Exobasidium vexans</i> AD4:- The consignment is free from <i>Cephaleuros parasiticus</i> , <i>Phomopsis theae</i> , & <i>Xanthomonas gortencovianum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant. (The material is to be of elite stock.)
<i>Campanula</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from Uredinales (Pine rusts).
<i>Cananga</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Capparis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Caragana</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Carissa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cassiope</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cedrela</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora cheoana</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceratonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceratopetalum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceropegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Chamaelaucium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cheiranthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Plasmiodiophora brassicae</i> & <i>Xanthomonas campestris</i> pv. <i>campestris</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Cicer</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ascochyta rabiei</i> , Beet curly top virus, <i>Verticillium albo-atrum</i> , & <i>Xanthomonas campestris</i> pv. <i>cassiae</i> did not occur on those plants or (ii) does not occur in the area of production concerned.

<i>Cistus</i>	AD4:- The consignment is free from <i>Phoma cisti</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Clematis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Clerodendrum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cneorum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Codonanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Coleus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cordyline</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> & <i>Phyllosticta dracaenae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Correa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cortaderia</i>	No additional declarations required. T6 = the consignment concerned was treated by an appropriate hot-water treatment.
<i>Costus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crescentia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crinodendron</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crossandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ctenanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cussonia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
Cycadales	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cytissus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Uromyces cytisi</i> , & <i>Uromyces genistae-tinctoriae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.

<i>Dahlia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> , & fungal wilt diseases did not occur on those plants or (ii) ds not occur in the area of production concerned.
<i>Dalbergia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Danae</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Daphne</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Dasyliiron</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Davidia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Delonix</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Delphinium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Beet curly top virus, & Tobacco ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Diascia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dieffenbachia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , <i>Phytophthora</i> spp. did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Dionaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dionysia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dioscorea</i>	AD4:- The consignment is free from <i>Goplana dioscorea</i> , & Yam mosaic virus.
<i>Dipladenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Doryanthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Draba</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dracaena</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> , & <i>Phyllosticta dracaenae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Echeveria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Edgeworthia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Elaeagnus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria galligena</i> , & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Epacris</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ephedra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Epipremnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Eremaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Erica</i>	AD4:- The consignment is free from <i>Phytophthora</i> spp., & <i>Verticillium</i> spp.
<i>Erigeron</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Eryngium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Accria peucedamum</i> , & <i>Aceria tulipae</i> .
<i>Erythrina</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Xanthomonas campestris</i> pv. <i>erythrinae</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Escallonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Euonymus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euonymus</i> variegation virus, & <i>Oidium euonymi-japonici</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Euphorbia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euphorbia</i> mosaic virus, & <i>Poinsettia</i> mosaic virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Euptelea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Eustoma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Exacum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Fittonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Flindersia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Hymenochaeta mongeotii</i> , did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Fremontia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Fuchsia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Gardenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Geraniaceae	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Pelargonium leaf curl virus, Tomato black ring virus, & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Gleditsia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Thyronectria denigrata</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Gmelina</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Grevillea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Halimocistus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hamamelis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> & <i>Polyporus gilvus</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Hebe</i>	T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hedera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Helianthemum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hemiandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide , T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hibbertia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Hosta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hoya</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hydrangea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Hydrangea ringspot virus, Tobacco ringspot virus, & Tomato ringspot virus, did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hyoscyamus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora hyoschyami</i> , & Tobacco rattle virus did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hypoestes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hyssopus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ilex</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Jatropha</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora jatrophiicola</i> , and <i>Phytophthora palmivora</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Kalanchoe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kalmia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Khaya</i>	AD4:- The consignment is free from <i>Xanthomonas campestris pv. khayae</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kolkwitzia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Arabis mosaic virus, Cherry leafroll virus, & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kunzea</i>	AD:- T1 = treated with a wide spectrum fungicide for control of <i>Phakopsora pachyrhizi</i> .
<i>Laburnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Tobacco ringspot virus, & <i>Uromyces genistae-tinctoriae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Lambertia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Euphorbia mosaic virus, & Poinsettia mosaic virus did not occur on those plants or (ii) do not occur in the area of production concerned.

<i>Lamium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Laurelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lavandula</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Leschenaultia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Limonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Liquidambar</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Liriodendron</i>	AD3:- The area of production is free from <i>Ceratocystis coerulescens</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lisianthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lonicera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Lychnis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lycopodium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lysimachia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lythrum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Magnolia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned, and <i>Nectria galligena</i> does not occur within 1 km of the area of production.
<i>Malva</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Abutilon mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. AD2:- :- New Zealand is free from Cotton leaf curl virus.
<i>Manilkara</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Medinilla</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Menianthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mesembryanthemum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora mesembryanthermi</i> did not occur or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Metrosideros</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mikania</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mimulus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora jacksonii</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Mimusops</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> , did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Mitella</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Monarda</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Moringa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Monstera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mussaenda</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Myrmecodia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nepenthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nepeta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nertera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Oenanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Oenothera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phymatotrichum omnivorum</i> , & <i>Verticillium albo-atrum</i> , did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Olearia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Onobrychis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Origanum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Patersonia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia convulata</i> , <i>Sclerotium tuliparum</i> , & Tobacco rattle virus did not occur on those plants or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> & <i>Ditylenchus dipsaci</i> .
<i>Peltiphyllum</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> & <i>Synchytrium endobioticum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Penstemon</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus, and <i>Phymatotrichum omnivorum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Peperomia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus, and Peperomia ringspot virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Persoonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Petasites</i>	AD:- T1 = treated with a wide spectrum fungicide for control of <i>Puccinia</i> spp. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Philadelphus</i>	AD4:- The consignment is free from Uredinales. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Philodendron</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Phlomis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Phlox</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pyrenochaeta phlogina</i> , Tomato aspermy virus, & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Phormium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Phormium yellowing mycoplasma, <i>Uredo phormii</i> , & <i>Xanthomonas campestris</i> pv. <i>phormicola</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Phylloglossum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Physostegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Phyteuma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Pimpinella</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Puccinia cari-bistortea</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD:- T1 = treated with a wide spectrum fungicide for control of <i>Puccinia cari-bistortea</i> . T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pisonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pittosporum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Platanus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ceratocystis fimbriata f.sp. platani</i> , & <i>Gnomonia platani</i> (syn. <i>G. veneta</i>), did not occur on those plants or (ii) do not occur in the area of production concerned. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Plectranthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Plumeria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polemonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polygonum</i>	AD4:- The consignment is free from <i>Uredo phormii</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Polypodiaceae	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polyscias</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Pothos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pouteria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Preslia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Prostanthera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Prunella</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Psilotum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pulmonaria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pyrostegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Rhamnus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria cinnabarina</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rhododendron</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Rhodohypoxis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rhoeo</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rivinia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rosmarinus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ruellia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas solanacearum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Rumex</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ruscus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Salix</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Physalospora miyabeana</i> , <i>Polaccia saliciperda</i> , <i>Pseudomonas saliciperda</i> , & Virus chlorosis did not occur on those plants or (ii) do not occur in the area of production concerned, and <i>Nectria galligena</i> did not occur within 1 Km of the place of production. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Salvia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phymatotrichum omnivorum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sambucus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus, Cherry leafroll virus & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Sansevieria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sarcocephalus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sarracenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Saxifraga</i>	AD4:- The consignment is free from Uredinales. AD:- T1 = treated with a wide spectrum fungicide for control of Uredinales.

<i>Scaevola</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Schefflera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Schizaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Scindapsus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Simmondsia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Soleirolia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sonerila</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Spathiphyllum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stachys</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stapelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Statice</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stephanotis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysanthemum stunt viroid</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Stereospermum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Symphoricarpos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Syngonium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Tabebuia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Talinum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Telopia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Thunbergia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tilia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Tmesipterus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tolmiea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Trachelium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tradescantia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tryphonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Veronica</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Verticordia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Viburnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Weigela</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus Cherry leafroll virus & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Wisteria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Wistringia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Xanthosoma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Yucca</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria obtusifolia</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Zizyphus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora zizyphi-vulgaris</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

4.4.2 Bulbs/Corms/Rhizomes/Tubers etc for propagation

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Bulbs/Corms/ Rhizomes /Tubers etc require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Aconitum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Agapanthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Aglaonema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Alstroemeria</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Amaryllidaceae (excluding species mentioned elsewhere)	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Pratylenchus scribneri</i> , <i>Sclerotium tuliparum</i> , & <i>Steneotarsonemus laticeps</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Androsace</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Aquilegia</i>	AD3:- The area of production is free from Beet curly top virus, Cucumber mosaic virus, <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Asparagus</i>	AD2:- New Zealand is free from Asparagus virus. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Astilbe</i>	AD2:- New Zealand is free from Asparagus virus. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Begonia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Xanthomonas campestris pv. begoniae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Beilis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Caladium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , & <i>Sclerotium tuliparum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cephalotus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , <i>Sclerotium tuliparum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Clivia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> <i>Pratylenchus scribneri</i> <i>Sclerotium tuliparum</i> , <i>Steneotarsonemus laticeps</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Colchicum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotium tuliparum</i> , <i>Urocystis cepulae</i> (Syn. <i>U. colchici</i>) did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Cyclamen</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ramularia cyclaminicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cynara</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Artichoke curly dwarf virus, Artichoke mosaic virus, Artichoke mottle crinkle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> and <i>Rotylenchus reniformis</i> AD11:- The consignment is free from soil or growth mediums.
<i>Dahlia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> & fungal wilt diseases did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , and <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Darlingtonia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Dicentra</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Eryngium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Accria peucedanum</i> , <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , and <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Gloriosa</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Heliotropium</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Hyacinthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , <i>Sclerotium tuliparum</i> , & <i>Xanthomonas campestris pv. hyacinthi</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Incarvillea</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Iridaceae (excluding species mentioned elsewhere)	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Drechslera iridis</i> , <i>Sclerotinia bulborum</i> , <i>Sclerotium tuliparum</i> , & Tobacco rattle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Lewisia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Liatis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Liliaceae (excluding species mentioned elsewhere)	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Lily mosaic virus, Lily rosette virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Macropidia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , & <i>Sclerotium tuliparum</i> , did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from Lily mosaic virus, Lily rosette virus, <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Marantaceae	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus, & <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Mirabilis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Mitella</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Myrmecodia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Oxalis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Paeonia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Panax</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Colletotrichum panicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Phlomis</i>	AD3:- The area of production is free from <i>Phomopsis vexans</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Phytax</i>	AD3:- The area of production is free from <i>Phomopsis vexans</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Sandersonia</i>	Refer to requirements Liliaceae.
<i>Sarracenia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Strelitzia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus, Banana bunchy top virus, <i>Erwinia carotovora</i> pv. <i>musae</i> , & <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Mycosphaerella fijiensis</i> , <i>Phymatotrichum omnivorum</i> , <i>Pseudomonas solanacearum</i> <i>Synchytrium endobioticum</i> , & <i>Xanthomonas campestris</i> pv. <i>celebensis</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Radopholus citrophilus</i> , & <i>Radopholus similis</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Tacca</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Thalictrum</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Zantedeschia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , & <i>Phytophthora</i> spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

4.4.3 Whole Plants

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Whole Plants require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below.

Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

Plants must be **glasshouse grown** and not older than 10 weeks and/or exceeding 200mm in height. All plants grown in the open ground (i.e. not within a glasshouse) require a phytosanitary import permit.

<i>Abelia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (E. pomi), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
---------------	---

<i>Acalypha</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Acanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Achillea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Adenia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Adenium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Aglaonema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Ajuga</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Albertia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Alocasia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Aloe</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Polyporus sanguineus</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Alstroemeria</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Amaranthaceae</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , Beet curly top virus, Cucumber mosaic virus, <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> , Tobacco ringspot virus, Tomato black ring virus, & Tomato ringspot virus. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Amaryllidaceae</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Androsace</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Anigozanthos</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Anthurium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Radopholus citrophilis</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Aphelandra</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Aralia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Arbutus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Ardisia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Arisaema</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Armeria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Artocarpus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Koleroga noxia</i> & <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Asparagus</i>	AD2:- New Zealand is free from Asparagus virus. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Astartea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Azalea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomya rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Baeckea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ballota</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Beaucarnea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bergenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bignonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Botrychium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bougainvillea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned, <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bouvardia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Brachychiton</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned, <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Brighamia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Buddleia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Calathea</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Calceolaria</i>	AD1:- (i) as appears from inspections during the active growth of that <i>Phytophthora</i> spp. did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Callistemon</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Camellia</i>	AD2:- New Zealand is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , and <i>Exobasidium vexans</i> . AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & <i>Xanthomonas grolencovianum</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Campanula</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & Uredinales. AD6:- The plants comprising the consignment were rooted and grown in sterilised media and were packed in sterilised media.
<i>Campsis</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Cananga</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Capparis</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Carissa</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Caulophyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ceanothus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ceratopetalum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Chamaelaucium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Cibotium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Clerodendron</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Clematis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cleyera</i>	AD2:- New Zealand is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , and <i>Exobasidium vexans</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Cephaluros parasiticus</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phomopsis theae</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & <i>Xanthomonas grolencovianum</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Codonanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Coleus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Columnea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Coprosma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Crossandra</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ctenanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cussonia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cyathea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Cycadales	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Cyclamen</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ramularia cyclaminicola</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dahlia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> & fungal wilt diseases did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dampiera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Danae</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dasylyrion</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Delphinium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Beet curly top virus, & Tobacco ringspot virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Dendranthema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Chrysanthemum stunt virus, <i>Didymella chrysanthemi</i> , <i>Erwinia chrysanthemi</i> , <i>Puccinia horiana</i> , & Tomato aspermy virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dianthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, <i>Cacoecimorpha pronubana</i> , Carnation etched ring virus, Carnation necrotic fleck virus, Carnation streak virus, <i>Erwinia chrysanthemi</i> pv. <i>dianthi</i> , Pelargonium leaf curl virus, <i>Pseudomonas caryophylli</i> , ring virus, Carnation necrotic fleck virus, Carnation streak virus, <i>Erwinia chrysanthemi</i> pv. <i>dianthi</i> , Pelargonium leaf curl virus, <i>Pseudomonas caryophylli</i> , & <i>Ustilago violacea</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted and grown in sterilised media. and were packed in sterilised media.
<i>Diascia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dicksonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dieffenbachia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , & <i>Phytophthora</i> spp., did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Dipladenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Draba</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dracaena</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> & <i>Phyllosticta dracaenae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Echeveria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Epipremnum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Erica</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora</i> spp., & <i>Verticillium</i> spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Erigeron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Eryngium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Euonymus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euonymus</i> variegation virus., & <i>Oidium euonymi-japonici</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Euphorbia</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD8:- Culture or tissue culture transplants were obtained from mother plants that were practically free of <i>Euphorbia</i> mosaic virus and <i>Poinsettia</i> mosaic virus.
<i>Eustoma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Exacum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Fatshedera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Fittonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Fuchsia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Gardenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Genista</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Geogenanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Geranium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Pelargonium leaf curl virus, Tomato black ring virus, <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Gerbera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora cryptogea</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Grevillea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Gypsophila</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hamamelis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> & <i>Polyporus gilvus</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Hebe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Hedera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Helichrysum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Helminthostachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hibbertia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hosta</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hoya</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Hydrangea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Hydrangea ringspot virus</i> , <i>Tobacco ringspot virus</i> & <i>Tomato ringspot virus</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hypoestes</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ilex</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ixora</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Cercospora ixorae</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Jatropha</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora jatrophiicola</i> & <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Kalanchoe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Kalmia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lambertia* AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD8:- culture or tissue culture transplants were obtained from mother plants that were practically free of Euphorbia mosaic virus and Poinsettia mosaic virus.
- Lamium* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
- Lavandula* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Leschenaultia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Limonium* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Lisianthus* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lonicera* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Verticillium albo-atrum* did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lychnis* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lycopodium* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lysimachia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Lythrum* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Macropidia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> & <i>Sclerotium tuliparum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , Lily mosaic virus, Lily rosette virus, <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Magnolia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria galligena</i> , <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Marantaceae	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus & <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Medinilla	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Menianthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Metrosideros</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Mimulus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora jacksonii</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Mitella</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Monarda</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Monstera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Mussaenda</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Myrmecodia</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nematanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Nepenthes</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nertera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nolina</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oenanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oenothera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ophioglossum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Origanum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Osmanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oxalis</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Ennomos subsignarius</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pachypodium</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pachystachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pandanus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cadang-cadang viroid did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Peltiphyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Penstemon</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Peperomia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus & Peperomia ringspot virus did not occur on those plants; or those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Petasites</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Puccinia</i> spp., <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Philodendron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Phlox</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pyrenochaeta phlogina</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Phormium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phormium</i> yellowing mycoplasma, <i>Uredo phormii</i> & <i>Xanthomonas campestris</i> pv. <i>phormicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Physostegia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Phyteuma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pisonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pittosporum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Polypodiaceae	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Polyscias</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Preslia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Primula</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Prunella</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pseuderanthemum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Psilotum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Pulmonaria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Punica</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sphaceloma punicae</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pyrenacantha</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pyrostegia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Rhadinophora</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> & <i>Phytophthora</i> spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Rhaphidiphora</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> & <i>Phytophthora</i> spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Rhododendron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Rheoe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ruellia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas solanacearum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ruscus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Salix</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Physalospora miyabeana</i> , <i>Polaccia saliciperda</i> , <i>Pseudomonas saliciperda</i> , & Virus chlorosis, did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Salvia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Sambucus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus, Cherry leafroll virus & Tobacco ringspot virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted and grown in sterilised media. and were packed in sterilised media.
<i>Sansevieria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Saxifraga</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Schefflera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Schismatoglottis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Schizaea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Scindapsus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Selaginella</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Simmondsia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Soleirolia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Sonerilla</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Spathiphyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Statice</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stephanotis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysanthemum stunt viroid</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stereospermum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Strelitzia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Abaca mosaic virus</i> , <i>Erwinia carotovora</i> pv. <i>musae</i> & <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Mycosphaerella fijiensis</i> , <i>Pseudomonas solanacearum</i> , & <i>Xanthomonas campestris</i> pv. <i>celebensis</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Syngonium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Talinum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Thalictrum</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tibouchina</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Thymus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tmesipterus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tolmiea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Trachelium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tradescantia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Verbascum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Veronica</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Viburnum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Wisteria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Xanthosoma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Yucca</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria obtusispora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

4.4.4 Tissue Culture *in vitro*

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Tissue Culture *in vitro* require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional

declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Achillea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Actinidia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aglaonema</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Alocasia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Amaranthaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Cucumber mosaic virus, Tobacco ringspot virus, Tomato black ring virus & Tomato ringspot virus.
Amaryllidaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Anigozanthos</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Anthurium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aphelandra</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aralia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ardisia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Asparagus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Asparagus virus.
<i>Aster</i>	AD4:- The consignment is free Uredinales (Pine rusts). AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, Tobacco rattle virus, & Tomato aspermy virus.
<i>Beaucarnea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Begonia</i> (excluding bulbous varieties)	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Botrychium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Bouvardia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Brachychiton</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Bromeliaceae (excluding <i>Ananas</i> spp.)	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Calathea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Campanula</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Cucumber mosaic virus,
<i>Ceropegia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Chamaelaucium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Clematis</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Clerodendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Codonanthe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Columnnea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cordyline</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Crossandra</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ctenanthe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cupressus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cussonia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dahlia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dendranthema</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, Chrysanthemum stunt viroid, & Tomato aspermy virus.

<i>Dianthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Carnation etched ring virus, Carnation necrotic fleck virus, Carnation streak virus, & Pelargonium leaf curl virus.
<i>Dieffenbachia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dipladenia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Draba</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dracaena</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Echeveria</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Erica</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Euphorbia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Euphorbia mosaic virus & Poinsettia mosaic virus.
<i>Fuchsia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Alfalfa dwarf mosaic virus.
<i>Geogenanthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Geranium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Pelargonium leaf curl virus, & Tomato black ring virus.
<i>Gerbera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Grevillea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Gypsophila</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hedera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Alfalfa dwarf mosaic virus.

<i>Heliconia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus & Banana bunchy top virus.
<i>Helminthostachys</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hibiscus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, & Cotton leaf curl virus.
<i>Hoya</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hydrangea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Hydrangea ringspot virus, Tobacco ringspot virus, & Tomato ringspot virus.
<i>Kalanchoe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Kalmia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lambertia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Euphorbia mosaic virus & Poinsettia mosaic virus.
<i>Limonium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lisianthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lycopodium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lysimachia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Marantaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus, & <i>Rosselinia bunodes</i> .
<i>Monstera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Nematanthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Nertera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ophioglossum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Ophiopogon</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Orchidaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Pachypodium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Passiflora</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Penstemon</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Cucumber mosaic virus.
<i>Philodendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Phlox</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Tomato aspermy virus.
<i>Phyloglossum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Phyteuma</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Pisonia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Polypodiaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Polyscias</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Psilotum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Rhododendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ruscus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Sansevieria</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Saxifraga</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Schizaea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Scindapsus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Simmondsia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Solidaster</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, & Tobacco rattle virus.
<i>Sonerilla</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Spathiphyllum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Stephanotis</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Chrysanthemum stunt viroid.
<i>Strelitzia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus & Banana bunchy top virus.
<i>Syngonium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Thalictrum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tibouchina</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tmesipterus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tradescantia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Yucca</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

4.5 Seeds (Grain)/Nuts

4.5.1 Seed for Sowing

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Seed for Sowing require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Abrus</i>	No additional declarations required.
<i>Abutilon</i>	No additional declarations required.
<i>Acaena</i>	Phytosanitary certificate not required.
<i>Acanthus</i>	Phytosanitary certificate not required.

<i>Achillea</i>	Phytosanitary certificate not required.
<i>Achimenes</i>	Phytosanitary certificate not required.
<i>Aciphylla</i>	Phytosanitary certificate not required.
<i>Aconitum</i>	Phytosanitary certificate not required.
<i>Acokanthera</i>	No additional declarations required.
<i>Acrocarpus</i>	No additional declarations required.
<i>Actaea</i>	Phytosanitary certificate not required.
<i>Actinidia</i>	Phytosanitary certificate not required.
<i>Actinotus</i>	Phytosanitary certificate not required.
<i>Adansonia</i>	No additional declarations required.
<i>Adenophora</i>	Phytosanitary certificate not required.
<i>Adenostyles</i>	Phytosanitary certificate not required.
<i>Adenanthos</i>	No additional declarations required.
<i>Adenium</i>	No additional declarations required.
<i>Aethionema</i>	Phytosanitary certificate not required.
<i>Aetoxicon</i>	No additional declarations required.
<i>Agastache</i>	Phytosanitary certificate not required.
<i>Agathis</i>	No additional declarations required.
<i>Agave</i>	No additional declarations required.
<i>Ageratum</i>	Phytosanitary certificate not required.
<i>Aglaonema</i>	Phytosanitary certificate not required.
<i>Agrostemma</i>	Phytosanitary certificate not required.
<i>Agonis</i>	No additional declarations required.
<i>Akebia</i>	No additional declarations required.
<i>Alberta</i>	No additional declarations required.
<i>Alcea</i>	Phytosanitary certificate not required.
<i>Alectryon</i>	No additional declarations required.
<i>Aleurites</i>	No additional declarations required.
<i>Alliaria</i>	Phytosanitary certificate not required.
<i>Alloteropsis</i>	No additional declarations required.
<i>Alluaudia</i>	No additional declarations required.
<i>Alnus</i>	No additional declarations required.

<i>Alocasia</i>	No additional declarations required.
<i>Aloe</i>	No additional declarations required.
<i>Aloysia</i>	No additional declarations required.
<i>Alseuosmia</i>	Phytosanitary certificate not required.
<i>Alstroemeria</i>	Phytosanitary certificate not required.
<i>Althaea</i>	Phytosanitary certificate not required.
<i>Alyssoides</i>	Phytosanitary certificate not required.
<i>Alyssum</i>	Phytosanitary certificate not required.
<i>Amaranthus</i>	Phytosanitary certificate not required.
<i>Amaryllis</i>	Phytosanitary certificate not required.
<i>Ammi</i>	Phytosanitary certificate not required.
<i>Ammobium</i>	Phytosanitary certificate not required.
<i>Amorpha</i>	No additional declarations required.
<i>Amorphophallus</i>	Phytosanitary certificate not required.
<i>Amphicarpaea</i>	No additional declarations required.
<i>Amsonia</i>	Phytosanitary certificate not required.
<i>Anacyclus</i>	Phytosanitary certificate not required.
<i>Ananas</i>	Phytosanitary certificate not required.
<i>Anaphalis</i>	Phytosanitary certificate not required.
<i>Anchusa</i>	Phytosanitary certificate not required.
<i>Angophora</i>	No additional declarations required.
<i>Anemone</i>	Phytosanitary certificate not required.
<i>Anethum</i>	Phytosanitary certificate not required.
<i>Angelica</i>	Phytosanitary certificate not required.
<i>Anigozanthos</i>	Phytosanitary certificate not required.
<i>Anthocleista</i>	No additional declarations required.
<i>Antennaria</i>	Phytosanitary certificate not required.
<i>Anthemis</i>	Phytosanitary certificate not required.
<i>Antholyza</i>	Phytosanitary certificate not required.
<i>Anthriscus</i>	Phytosanitary certificate not required.
<i>Anthurium</i>	Phytosanitary certificate not required.
<i>Anthyllis</i>	Phytosanitary certificate not required.

<i>Antidesma</i>	No additional declarations required.
<i>Antigonon</i>	No additional declarations required.
<i>Antirrhinum</i>	Phytosanitary certificate not required.
<i>Aphelandra</i>	Phytosanitary certificate not required.
<i>Apios</i>	Phytosanitary certificate not required.
<i>Apocynum</i>	Phytosanitary certificate not required.
<i>Aquilegia</i>	Phytosanitary certificate not required.
<i>Aralia</i>	No additional declarations required.
<i>Arbutus</i>	No additional declarations required.
<i>Arctium</i>	Phytosanitary certificate not required.
<i>Arctotis</i>	Phytosanitary certificate not required.
<i>Arctostaphylos</i>	No additional declarations required.
<i>Ardisa</i>	No additional declarations required.
<i>Argyroderma</i>	Phytosanitary certificate not required.
<i>Arisaema</i>	No additional declarations required.
<i>Aristolochia</i>	No additional declarations required.
<i>Armeria</i>	Phytosanitary certificate not required.
<i>Artemisia</i>	Phytosanitary certificate not required.
<i>Artocarpus</i>	No additional declarations required.
<i>Arum</i>	No additional declarations required.
<i>Asarina</i>	Phytosanitary certificate not required.
<i>Asclepias</i>	Phytosanitary certificate not required.
<i>Asimina</i>	No additional declarations required.
<i>Astartea</i>	No additional declarations required.
<i>Aster</i>	Phytosanitary certificate not required.
<i>Astragalus</i>	Phytosanitary certificate not required.
<i>Astrantia</i>	Phytosanitary certificate not required.
<i>Asyneuma</i>	Phytosanitary certificate not required.
<i>Atalaya</i>	No additional declarations required.
<i>Atriplex</i>	No additional declarations required.
<i>Aubrieta</i>	Phytosanitary certificate not required.
<i>Aurinia</i>	Phytosanitary certificate not required.

<i>Axonopus</i>	Phytosanitary certificate not required.
<i>Azadirachta</i>	No additional declarations required.
<i>Azanza</i>	No additional declarations required.
<i>Azara</i>	No additional declarations required.
<i>Baeckea</i>	No additional declarations required.
<i>Balanites</i>	No additional declarations required.
<i>Ballota</i>	Phytosanitary certificate not required.
<i>Balsamita</i>	Phytosanitary certificate not required.
<i>Banksia</i>	No additional declarations required.
<i>Baptisia</i>	Phytosanitary certificate not required.
<i>Basilicum</i>	Phytosanitary certificate not required.
<i>Bassia</i>	Phytosanitary certificate not required.
<i>Bauhinia</i>	No additional declarations required.
<i>Beaucarnea</i>	No additional declarations required.
<i>Beaufortia</i>	No additional declarations required.
<i>Beaumontia</i>	No additional declarations required.
<i>Begonia</i>	Phytosanitary certificate not required.
<i>Bergenia</i>	Phytosanitary certificate not required.
<i>Berlandiera</i>	Phytosanitary certificate not required.
<i>Beschorneria</i>	Phytosanitary certificate not required.
<i>Bignonia</i>	No additional declarations required.
<i>Bischofia</i>	No additional declarations required.
<i>Biserrula</i>	No additional declarations required.
<i>Boltonia</i>	Phytosanitary certificate not required.
<i>Borago</i>	Phytosanitary certificate not required.
<i>Bossiaea</i>	No additional declarations required.
<i>Botrychium</i>	No additional declarations required.
<i>Bougainvillea</i>	No additional declarations required.
<i>Bouvardia</i>	Phytosanitary certificate not required.
<i>Brachychiton</i>	No additional declarations required.
<i>Brachycome</i>	Phytosanitary certificate not required.
<i>Brachylaena</i>	No additional declarations required.

<i>Brachysema</i>	Phytosanitary certificate not required.
<i>Brachystelma</i>	Phytosanitary certificate not required.
<i>Brachystemma</i>	No additional declarations required.
<i>Brachystegia</i>	Phytosanitary certificate not required.
<i>Bridelia</i>	No additional declarations required.
<i>Brighamia</i>	No additional declarations required.
<i>Brizia</i>	No additional declarations required.
<i>Bromelia</i>	Phytosanitary certificate not required.
<i>Brownea</i>	No additional declarations required.
<i>Browallia</i>	Phytosanitary certificate not required.
<i>Brugmansia</i>	Phytosanitary certificate not required.
<i>Brunfelsia</i>	No additional declarations required.
<i>Buckinghamia</i>	No additional declarations required.
<i>Buddleia</i>	No additional declarations required.
<i>Bupleurum</i>	No additional declarations required.
<i>Buphthalmum</i>	Phytosanitary certificate not required.
<i>Burchellia</i>	No additional declarations required.
<i>Bursaria</i>	No additional declarations required.
<i>Bursera</i>	No additional declarations required.
<i>Byblis</i>	Phytosanitary certificate not required.
<i>Cajanus</i>	No additional declarations required.
<i>Caladium</i>	No additional declarations required.
<i>Calamintha</i>	Phytosanitary certificate not required.
<i>Calamus</i>	No additional declarations required.
<i>Calathea</i>	Phytosanitary certificate not required.
<i>Calceolaria</i>	Phytosanitary certificate not required.
<i>Calendula</i>	Phytosanitary certificate not required.
<i>Calliandra</i>	No additional declarations required.
<i>Callistemon</i>	No additional declarations required.
<i>Callistephus</i>	Phytosanitary certificate not required.
<i>Callitris</i>	No additional declarations required.
<i>Calocephalus</i>	Phytosanitary certificate not required.

<i>Calothamnus</i>	No additional declarations required.
<i>Calpurnia</i>	No additional declarations required.
<i>Caltha</i>	Phytosanitary certificate not required.
<i>Calycanthus</i>	Phytosanitary certificate not required.
<i>Campanula</i>	Phytosanitary certificate not required.
<i>Cananga</i>	No additional declarations required.
<i>Canavalia</i>	Phytosanitary certificate not required.
<i>Canna</i>	Phytosanitary certificate not required.
<i>Canthium</i>	No additional declarations required.
<i>Capparis</i>	No additional declarations required.
<i>Carangana</i>	No additional declarations required.
<i>Caralluma</i>	Phytosanitary certificate not required.
<i>Carissa</i>	No additional declarations required.
<i>Carlina</i>	Phytosanitary certificate not required.
<i>Carludovica</i>	Phytosanitary certificate not required.
<i>Carpanthea</i>	Phytosanitary certificate not required.
<i>Carpinus</i>	No additional declarations required.
<i>Carpodetus</i>	No additional declarations required.
<i>Carthamus</i>	Phytosanitary certificate not required.
<i>Caryocar</i>	No additional declarations required.
<i>Caryopteris</i>	No additional declarations required.
<i>Cassia</i>	No additional declarations required.
<i>Cassine</i>	Phytosanitary certificate not required.
<i>Cassiope</i>	No additional declarations required.
<i>Catanache</i>	Phytosanitary certificate not required.
<i>Catharanthus</i>	Phytosanitary certificate not required.
<i>Ceanothus</i>	No additional declarations required.
<i>Cecropia</i>	No additional declarations required.
<i>Cedrela</i>	No additional declarations required.
<i>Cedrus</i>	No additional declarations required.
<i>Celosia</i>	Phytosanitary certificate not required.
<i>Celtis</i>	No additional declarations required.

<i>Cenchrus</i>	No additional declarations required.
<i>Centaurea</i>	Phytosanitary certificate not required.
<i>Centranthus</i>	Phytosanitary certificate not required.
<i>Cephalaria</i>	Phytosanitary certificate not required.
<i>Cephalotaxus</i>	No additional declarations required.
<i>Cerastium</i>	Phytosanitary certificate not required.
<i>Ceratonia</i>	No additional declarations required.
<i>Ceratopetalum</i>	No additional declarations required.
<i>Cercidium</i>	No additional declarations required.
<i>Cerinth</i>	Phytosanitary certificate not required.
<i>Ceropegia</i>	Phytosanitary certificate not required.
<i>Cercis</i>	No additional declarations required.
<i>Chamaecytisus</i>	No additional declarations required.
<i>Chamaelaucium</i>	No additional declarations required.
<i>Cheiranthus</i>	Phytosanitary certificate not required.
<i>Chelone</i>	Phytosanitary certificate not required.
<i>Chionanthus</i>	No additional declarations required.
<i>Chionochoila</i>	No additional declarations required.
<i>Chloris</i>	No additional declarations required.
<i>Chorizema</i>	No additional declarations required.
<i>Chrysophyllum</i>	No additional declarations required.
<i>Chrysopsis</i>	Phytosanitary certificate not required.
<i>Chukrasia</i>	No additional declarations required.
<i>Cichorium</i>	No additional declarations required.
<i>Cimicifuga</i>	Phytosanitary certificate not required.
<i>Cineraria</i>	Phytosanitary certificate not required.
<i>Cinnamomum</i>	No additional declarations required.
<i>Cistus</i>	No additional declarations required.
<i>Cladanthus</i>	Phytosanitary certificate not required.
<i>Cladrastis</i>	No additional declarations required.
<i>Clarkia</i>	Phytosanitary certificate not required.
<i>Cleome</i>	Phytosanitary certificate not required.

<i>Clerodendrum</i>	No additional declarations required.
<i>Clianthus</i>	No additional declarations required.
<i>Clinopodium</i>	Phytosanitary certificate not required.
<i>Clitoria</i>	No additional declarations required.
<i>Clivia</i>	Phytosanitary certificate not required.
<i>Cneorum</i>	No additional declarations required.
<i>Cobaea</i>	No additional declarations required.
<i>Coccoloba</i>	No additional declarations required.
<i>Cochlearia</i>	Phytosanitary certificate not required.
<i>Cochlospermum</i>	No additional declarations required.
<i>Codiaeum</i>	No additional declarations required.
<i>Codonopsis</i>	Phytosanitary certificate not required.
<i>Cola</i>	No additional declarations required.
<i>Colchicum</i>	No additional declarations required.
<i>Coleus</i>	Phytosanitary certificate not required.
<i>Collinsia</i>	Phytosanitary certificate not required.
<i>Colubrina</i>	No additional declarations required.
<i>Colutea</i>	No additional declarations required.
<i>Colvillea</i>	No additional declarations required.
<i>Commiphora</i>	Phytosanitary certificate not required.
<i>Conophytum</i>	Phytosanitary certificate not required.
<i>Consolida</i>	Phytosanitary certificate not required.
<i>Convolvulus</i>	Phytosanitary certificate not required.
<i>Coprosma</i>	No additional declarations required.
<i>Cordyline</i>	No additional declarations required.
<i>Coriandrum</i>	Phytosanitary certificate not required.
<i>Corema</i>	No additional declarations required.
<i>Coriaria</i>	No additional declarations required.
<i>Coreopsis</i>	Phytosanitary certificate not required.
<i>Coronilla</i>	No additional declarations required.
<i>Cortaderia</i>	No additional declarations required.
<i>Corydalis</i>	No additional declarations required.

<i>Corynocarpus</i>	No additional declarations required.
<i>Cosmos</i>	Phytosanitary certificate not required.
<i>Cotinus</i>	No additional declarations required.
<i>Cotoneaster</i>	No additional declarations required.
<i>Crambe</i>	Phytosanitary certificate not required.
<i>Crepis</i>	Phytosanitary certificate not required.
<i>Crescentia</i>	Phytosanitary certificate not required.
<i>Crinodendron</i>	No additional declarations required.
<i>Crossandra</i>	Phytosanitary certificate not required.
<i>Croton</i>	No additional declarations required.
<i>Cryptotaenia</i>	Phytosanitary certificate not required.
<i>Cuminum</i>	Phytosanitary certificate not required.
<i>Cupaniopsis</i>	No additional declarations required.
<i>Cuphea</i>	Phytosanitary certificate not required.
<i>Curculigo</i>	Phytosanitary certificate not required.
<i>Curcuma</i>	Phytosanitary certificate not required.
<i>Cussonia</i>	No additional declarations required.
<i>Cyathea</i>	No additional declarations required.
Cycadales	No additional declarations required.
<i>Cyclamen</i>	Phytosanitary certificate not required.
<i>Cymbopogon</i>	No additional declarations required.
<i>Cynara</i>	Phytosanitary certificate not required.
<i>Cynodon</i>	No additional declarations required.
<i>Cynoglossum</i>	Phytosanitary certificate not required.
<i>Cytisus</i>	No additional declarations required.
<i>Dahlia</i>	Phytosanitary certificate not required.
<i>Dais</i>	No additional declarations required.
<i>Dalbergia</i>	No additional declarations required.
<i>Danae</i>	No additional declarations required.
<i>Darlingtonia</i>	Phytosanitary certificate not required.
<i>Daphne</i>	No additional declarations required.
<i>Darwinia</i>	No additional declarations required.

<i>Dasyilirion</i>	No additional declarations required.
<i>Davidia</i>	No additional declarations required.
<i>Delonix</i>	No additional declarations required.
<i>Diascia</i>	Phytosanitary certificate not required.
<i>Dialium</i>	No additional declarations required.
<i>Dicentra</i>	Phytosanitary certificate not required.
<i>Dichondra</i>	Phytosanitary certificate not required.
<i>Didelta</i>	Phytosanitary certificate not required.
<i>Didierea</i>	No additional declarations required.
<i>Didymaotus</i>	No additional declarations required.
<i>Dieffenbachia</i>	No additional declarations required.
<i>Dillenia</i>	No additional declarations required.
<i>Dimocarpus</i>	No additional declarations required.
<i>Dimorphotheca</i>	Phytosanitary certificate not required.
<i>Dionaea</i>	Phytosanitary certificate not required.
<i>Dioscorea</i>	Phytosanitary certificate not required.
<i>Dipsacus</i>	Phytosanitary certificate not required.
<i>Dipterocarpus</i>	No additional declarations required.
<i>Dischidia</i>	Phytosanitary certificate not required.
<i>Dodecatheon</i>	Phytosanitary certificate not required.
<i>Dodonaea</i>	No additional declarations required.
<i>Dombeya</i>	No additional declarations required.
<i>Doronicum</i>	Phytosanitary certificate not required.
<i>Dorstenia</i>	No additional declarations required.
<i>Doryanthes</i>	Phytosanitary certificate not required.
<i>Draba</i>	Phytosanitary certificate not required.
<i>Dracaena</i>	No additional declarations required.
<i>Dracocephalum</i>	Phytosanitary certificate not required.
<i>Drosera</i>	Phytosanitary certificate not required.
<i>Dryandra</i>	No additional declarations required.
<i>Dysoxylum</i>	No additional declarations required.
<i>Duvalia</i>	Phytosanitary certificate not required.

<i>Echeveria</i>	Phytosanitary certificate not required.
<i>Echinacea</i>	Phytosanitary certificate not required.
<i>Echinops</i>	Phytosanitary certificate not required.
<i>Echium</i>	Phytosanitary certificate not required.
<i>Ehretia</i>	No additional declarations required.
<i>Elacocarpus</i>	No additional declarations required.
<i>Elaeagnus</i>	No additional declarations required.
<i>Elettaria</i>	Phytosanitary certificate not required.
<i>Eleutherococcus</i>	No additional declarations required.
<i>Edgeworthia</i>	No additional declarations required.
<i>Enterolobium</i>	No additional declarations required.
<i>Epacris</i>	No additional declarations required.
<i>EphedraEpilobium</i>	No additional declarations required. Phytosanitary certificate not required.
<i>Eremophila</i>	No additional declarations required.
<i>Erigeron</i>	Phytosanitary certificate not required.
<i>Eriobotrya</i>	No additional declarations required.
<i>Eriogonum</i>	Phytosanitary certificate not required.
<i>Eriophyllum</i>	No additional declarations required.
<i>Eruca</i>	Phytosanitary certificate not required.
<i>Eryngium</i>	Phytosanitary certificate not required.
<i>Erysimum</i>	Phytosanitary certificate not required.
<i>Erythrina</i>	No additional declarations required.
<i>Escallonia</i>	No additional declarations required.
<i>Eschscholzia</i>	Phytosanitary certificate not required.
<i>Euclea</i>	No additional declarations required.
<i>Eucommia</i>	No additional declarations required.
<i>Eucryphia</i>	No additional declarations required.
<i>Eupatorium</i>	Phytosanitary certificate not required.
<i>Euphorbia</i>	No additional declarations required.
<i>Euryops</i>	Phytosanitary certificate not required.
<i>Eustoma</i>	Phytosanitary certificate not required.
<i>Euptelea</i>	No additional declarations required.

<i>Evolvulus</i>	Phytosanitary certificate not required.
<i>Exacum</i>	Phytosanitary certificate not required.
<i>Faucaria</i>	No additional declarations required.
<i>Felicia</i>	Phytosanitary certificate not required.
<i>Ficus</i>	No additional declarations required.
<i>Fittonia</i>	Phytosanitary certificate not required.
<i>Flacourtia</i>	No additional declarations required.
<i>Flindersia</i>	No additional declarations required.
<i>Foeniculum</i>	Phytosanitary certificate not required.
<i>Fremontodenron</i>	No additional declarations required.
<i>Freesia</i>	Phytosanitary certificate not required.
<i>Fuchsia</i>	Phytosanitary certificate not required.
<i>Fumaria</i>	Phytosanitary certificate not required.
<i>Furcraea</i>	No additional declarations required.
<i>Gahnia</i>	No additional declarations required.
<i>Gaillardia</i>	No additional declarations required.
<i>Galega</i>	Phytosanitary certificate not required.
<i>Galium</i>	Phytosanitary certificate not required.
<i>Galphimia</i>	No additional declarations required.
<i>Gardenia</i>	No additional declarations required.
<i>Garcinia</i>	No additional declarations required.
<i>Gaura</i>	Phytosanitary certificate not required.
<i>Gazania</i>	Phytosanitary certificate not required.
<i>Genista</i>	No additional declarations required.
<i>Gentiana</i>	Phytosanitary certificate not required.
<i>Geranium</i>	Phytosanitary certificate not required.
<i>Gerbera</i>	Phytosanitary certificate not required.
<i>Geum</i>	Phytosanitary certificate not required.
<i>Gilia</i>	Phytosanitary certificate not required.
<i>Ginkgo</i>	No additional declarations required.
<i>Gladiolus</i>	Phytosanitary certificate not required.
<i>Glaucium</i>	Phytosanitary certificate not required.

<i>Gleditsia</i>	No additional declarations required.
<i>Gliricidia</i>	No additional declarations required.
<i>Globularia</i>	Phytosanitary certificate not required.
<i>Gloxinia</i>	Phytosanitary certificate not required.
<i>Gmelina</i>	No additional declarations required.
<i>Godetia</i>	Phytosanitary certificate not required.
<i>Gomphrena</i>	Phytosanitary certificate not required.
<i>Goodenia</i>	No additional declarations required.
<i>Gossypium</i>	No additional declarations required. Must be acid de-linted.
<i>Gosswilerodendron</i>	No additional declarations required.
<i>Gouania</i>	No additional declarations required.
<i>Grevillea</i>	No additional declarations required.
<i>Greyia</i>	No additional declarations required.
<i>Grindelia</i>	Phytosanitary certificate not required.
<i>Griselina</i>	No additional declarations required.
<i>Gymnosporia</i>	No additional declarations required.
<i>Gypsophila</i>	Phytosanitary certificate not required.
<i>Haloxylon</i>	No additional declarations required.
<i>Hamamelis</i>	No additional declarations required.
<i>Hardenbergia</i>	No additional declarations required.
<i>Harungana</i>	No additional declarations required.
<i>Hebe</i>	No additional declarations required.
<i>Hedera</i>	Phytosanitary certificate not required.
<i>Hebenstretia</i>	No additional declarations required.
<i>Hedysarum</i>	No additional declarations required.
<i>Helenium</i>	Phytosanitary certificate not required.
<i>Helichrysum</i>	Phytosanitary certificate not required.
<i>Helinus</i>	No additional declarations required.
<i>Heliophila</i>	Phytosanitary certificate not required.
<i>Heliopsis</i>	Phytosanitary certificate not required.
<i>Heliocarpus</i>	No additional declarations required.
<i>Helleborus</i>	No additional declarations required.

<i>Helminthostachys</i>	No additional declarations required.
<i>Hemigenia</i>	No additional declarations required.
<i>Heliotropium</i>	Phytosanitary certificate not required.
<i>Helipterum</i>	Phytosanitary certificate not required.
<i>Hesperis</i>	Phytosanitary certificate not required.
<i>Heuchera</i>	Phytosanitary certificate not required.
<i>Hibbertia</i>	No additional declarations required.
<i>Hieracium</i>	Phytosanitary certificate not required.
<i>Hildegardia</i>	No additional declarations required.
<i>Hippophae</i>	No additional declarations required.
<i>Hoheria</i>	No additional declarations required.
<i>Hosta</i>	Phytosanitary certificate not required.
<i>Hovea</i>	No additional declarations required.
<i>Hovenia</i>	No additional declarations required.
<i>Hyacinthus</i>	Phytosanitary certificate not required.
<i>Hydrocotyle</i>	Phytosanitary certificate not required.
<i>Hymenachne</i>	No additional declarations required.
<i>Hymenantherum</i>	No additional declarations required.
<i>Hymenosporum</i>	No additional declarations required.
<i>Hyoscyamus</i>	Phytosanitary certificate not required.
<i>Hypocalymma</i>	No additional declarations required.
<i>Hypoestes</i>	Phytosanitary certificate not required.
<i>Hyssopus</i>	Phytosanitary certificate not required.
<i>Iberis</i>	Phytosanitary certificate not required.
<i>Illex</i>	No additional declarations required.
<i>Impatiens</i>	No additional declarations required.
<i>Incarvillea</i>	Phytosanitary certificate not required.
<i>Indigofera</i>	No additional declarations required.
<i>Inula</i>	Phytosanitary certificate not required.
<i>Isatis</i>	Phytosanitary certificate not required.
<i>Ixodia</i>	Phytosanitary certificate not required.
<i>Ixora</i>	No additional declarations required.

<i>Jasione</i>	Phytosanitary certificate not required.
<i>Jatropha</i>	No additional declarations required.
<i>Justica</i>	No additional declarations required.
<i>Jurinea</i>	Phytosanitary certificate not required.
<i>Kalanchoe</i>	Phytosanitary certificate not required.
<i>Kalmia</i>	No additional declarations required.
<i>Kennedia</i>	No additional declarations required.
<i>Khaya</i>	No additional declarations required.
<i>Kiggelaria</i>	No additional declarations required.
<i>Kirengeshoma</i>	No additional declarations required.
<i>Kirkia</i>	No additional declarations required.
<i>Kissenia</i>	No additional declarations required.
<i>Knautia</i>	Phytosanitary certificate not required.
<i>Knightsia</i>	No additional declarations required.
<i>Kniphofia</i>	Phytosanitary certificate not required.
<i>Kochia</i>	Phytosanitary certificate not required.
<i>Koelreuteria</i>	No additional declarations required.
<i>Kolkritzia</i>	No additional declarations required.
<i>Kunzea</i>	No additional declarations required.
<i>Laburnum</i>	No additional declarations required.
<i>Lachnostachys</i>	Phytosanitary certificate not required.
<i>Lagerstroemia</i>	No additional declarations required.
<i>Landolphia</i>	No additional declarations required.
<i>Langsdorffia</i>	No additional declarations required.
<i>Lansea</i>	No additional declarations required.
<i>Laportea</i>	No additional declarations required.
<i>Lasthenia</i>	Phytosanitary certificate not required.
<i>Lathyrus</i>	Phytosanitary certificate not required.
<i>Larix</i>	No additional declarations required.
<i>Laurelia</i>	No additional declarations required.
<i>Laurentia</i>	Phytosanitary certificate not required.
<i>Laurus</i>	No additional declarations required.

<i>Lavandula</i>	Phytosanitary certificate not required.
<i>Lavatera</i>	Phytosanitary certificate not required.
<i>Layia</i>	Phytosanitary certificate not required.
<i>Leonotis</i>	Phytosanitary certificate not required.
<i>Leontopodium</i>	Phytosanitary certificate not required.
<i>Lepidium</i>	Phytosanitary certificate not required.
<i>Lechenaultia</i>	No additional declarations required.
<i>Lespedeza</i>	Phytosanitary certificate not required.
<i>Lesquerella</i>	No additional declarations required.
<i>Leucanthemum</i>	Phytosanitary certificate not required.
<i>Leucopogon</i>	No additional declarations required.
<i>Leucothoe</i>	No additional declarations required.
<i>Levisticum</i>	Phytosanitary certificate not required.
<i>Lewisia</i>	Phytosanitary certificate not required.
<i>Liatris</i>	Phytosanitary certificate not required.
<i>Ligularia</i>	Phytosanitary certificate not required.
<i>Lilium</i>	Phytosanitary certificate not required.
<i>Limnanthes</i>	Phytosanitary certificate not required.
<i>Limonium</i>	Phytosanitary certificate not required.
<i>Linaria</i>	Phytosanitary certificate not required.
<i>Lindera</i>	No additional declarations required.
<i>Lisianthus</i>	Phytosanitary certificate not required.
<i>Liquidambar</i>	No additional declarations required.
<i>Litchi</i>	No additional declarations required.
<i>Lithops</i>	No additional declarations required.
<i>Lobelia</i>	No additional declarations required.
<i>Lobularia</i>	Phytosanitary certificate not required.
<i>Lomatium</i>	Phytosanitary certificate not required.
<i>Lonas</i>	Phytosanitary certificate not required.
<i>Lophiocarpus</i>	No additional declarations required.
<i>Lophotocarpus</i>	No additional declarations required.
<i>Lotus</i>	No additional declarations required.

<i>Luculia</i>	No additional declarations required.
<i>Luehea</i>	No additional declarations required.
<i>Lunaria</i>	Phytosanitary certificate not required.
<i>Luzula</i>	Phytosanitary certificate not required.
<i>Lychnis</i>	Phytosanitary certificate not required.
<i>Lycopodiella</i>	Phytosanitary certificate not required.
<i>Lycopodium</i>	Phytosanitary certificate not required.
<i>Lycopus</i>	Phytosanitary certificate not required.
<i>Lysichiton</i>	Phytosanitary certificate not required.
<i>Lysimachia</i>	Phytosanitary certificate not required.
<i>Lythrum</i>	Phytosanitary certificate not required.
<i>Macfadyena</i>	No additional declarations required.
<i>Macarthuria</i>	Phytosanitary certificate not required.
<i>Macleaya</i>	Phytosanitary certificate not required.
<i>Macropidia</i>	Phytosanitary certificate not required.
<i>Macropiper</i>	No additional declarations required.
<i>Macroptilium</i>	No additional declarations required.
<i>Maesa</i>	No additional declarations required.
<i>Magnolia</i>	No additional declarations required.
<i>Malcolmia</i>	Phytosanitary certificate not required.
<i>Malva</i>	Phytosanitary certificate not required.
<i>Mammea</i>	No additional declarations required.
<i>Mandevilla</i>	No additional declarations required.
<i>Manilkara</i>	No additional declarations required.
<i>Maranta</i>	Phytosanitary certificate not required.
<i>Markhamia</i>	No additional declarations required.
<i>Marrubium</i>	No additional declarations required.
<i>Matricaria</i>	Phytosanitary certificate not required.
<i>Matthiola</i>	Phytosanitary certificate not required.
<i>Medinilla</i>	No additional declarations required.
<i>Meconopsis</i>	Phytosanitary certificate not required.
<i>Melaleuca</i>	No additional declarations required.

<i>Medusagyne</i>	Phytosanitary certificate not required.
<i>Melampodium</i>	Phytosanitary certificate not required.
<i>Melianthus</i>	No additional declarations required.
<i>Melissa</i>	Phytosanitary certificate not required.
<i>Memecylon</i>	No additional declarations required.
<i>Mentzelia</i>	No additional declarations required.
<i>Menyanthes</i>	Phytosanitary certificate not required.
<i>Mertensia</i>	Phytosanitary certificate not required.
<i>Meryta</i>	No additional declarations required.
<i>Mesembryanthemum</i>	Phytosanitary certificate not required.
<i>Metrosideros</i>	No additional declarations required.
<i>Meum</i>	No additional declarations required.
<i>Miconia</i>	No additional declarations required.
<i>Mikania</i>	No additional declarations required.
<i>Millettia</i>	No additional declarations required.
<i>Mimosa</i>	No additional declarations required.
<i>Mimulus</i>	Phytosanitary certificate not required.
<i>Mimusops</i>	No additional declarations required.
<i>Mina</i>	Phytosanitary certificate not required.
<i>Momordica</i>	No additional declarations required.
<i>Mirabilis</i>	Phytosanitary certificate not required.
<i>Moluccella</i>	Phytosanitary certificate not required.
<i>Monarda</i>	Phytosanitary certificate not required.
<i>Monstera</i>	Phytosanitary certificate not required.
<i>Morina</i>	Phytosanitary certificate not required.
<i>Moringa</i>	No additional declarations required.
<i>Mundulea</i>	Phytosanitary certificate not required.
<i>Mucuna</i>	No additional declarations required.
<i>Myosotidium</i>	Phytosanitary certificate not required.
<i>Myristica</i>	No additional declarations required.
<i>Myrrhis</i>	Phytosanitary certificate not required.

<i>Myrsine</i>	No additional declarations required.
<i>Myrtus</i>	No additional declarations required.
<i>Nandina</i>	Phytosanitary certificate not required.
<i>Nasturtium</i>	Phytosanitary certificate not required.
<i>Nemesia</i>	Phytosanitary certificate not required.
<i>Nemophila</i>	Phytosanitary certificate not required.
<i>Nepenthes</i>	Phytosanitary certificate not required.
<i>Nerium</i>	No additional declarations required.
<i>Nepeta</i>	Phytosanitary certificate not required.
<i>Nertera</i>	Phytosanitary certificate not required.
<i>Newtonia</i>	Phytosanitary certificate not required.
<i>Nierembergia</i>	Phytosanitary certificate not required.
<i>Nigella</i>	Phytosanitary certificate not required.
<i>Nolana</i>	Phytosanitary certificate not required.
<i>Nuxia</i>	No additional declarations required.
<i>Nuytsia</i>	No additional declarations required.
<i>Nymphaea</i>	No additional declarations required.
<i>Nyssa</i>	No additional declarations required.
<i>Ocimum</i>	Phytosanitary certificate not required.
<i>Oenothera</i>	Phytosanitary certificate not required.
<i>Olearia</i>	Phytosanitary certificate not required.
<i>Oncoba</i>	No additional declarations required.
<i>Ongokea</i>	No additional declarations required.
<i>Onobrychis</i>	No additional declarations required.
<i>Operculicarya</i>	No additional declarations required.
<i>Ophioglossum</i>	No additional declarations required.
Orchidaceae	No additional declarations required.
<i>Origanum</i>	Phytosanitary certificate not required.
<i>Ornithopus</i>	Phytosanitary certificate not required.
<i>Osteospermum</i>	Phytosanitary certificate not required.
<i>Ostrya</i>	No additional declarations required.
<i>Osyrius</i>	No additional declarations required.

<i>Oxypetalum</i>	Phytosanitary certificate not required.
<i>Pachypodium</i>	Phytosanitary certificate not required.
<i>Paeonia</i>	Phytosanitary certificate not required.
<i>Parnassia</i>	Phytosanitary certificate not required.
<i>Parinari</i>	No additional declarations required.
<i>Parkinsonia</i>	No additional declarations required.
<i>Parthenium</i>	Phytosanitary certificate not required.
<i>Pastinaca</i>	Phytosanitary certificate not required.
<i>Paulownia</i>	No additional declarations required.
<i>Pavetta</i>	No additional declarations required.
<i>Peganum</i>	No additional declarations required.
<i>Peltiphyllum</i>	Phytosanitary certificate not required.
<i>Peltophorum</i>	No additional declarations required.
<i>Pentas</i>	No additional declarations required.
<i>Penstemon</i>	Phytosanitary certificate not required.
<i>Peperomia</i>	Phytosanitary certificate not required.
<i>Perilla</i>	Phytosanitary certificate not required.
<i>Perovskia</i>	No additional declarations required.
<i>Persoonia</i>	No additional declarations required.
<i>Petalostemon</i>	Phytosanitary certificate not required.
<i>Petasites</i>	Phytosanitary certificate not required.
<i>Petrophila</i>	No additional declarations required.
<i>Petroselinum</i>	Phytosanitary certificate not required.
<i>Petunia</i>	Phytosanitary certificate not required.
<i>Peucedanum</i>	Phytosanitary certificate not required.
<i>Phacelia</i>	Phytosanitary certificate not required.
<i>Phalaris</i>	No additional declarations required.
<i>Philodendron</i>	Phytosanitary certificate not required.
<i>Phleum</i>	No additional declarations required.
<i>Phlomis</i>	No additional declarations required.
<i>Phlox</i>	No additional declarations required.
<i>Phormium</i>	No additional declarations required.

<i>Phygelius</i>	No additional declarations required.
<i>Phylloglossum</i>	Phytosanitary certificate not required.
<i>Phyllanthus</i>	No additional declarations required.
<i>Phyllocladus</i>	No additional declarations required.
<i>Physostegia</i>	Phytosanitary certificate not required.
<i>Phyteuma</i>	Phytosanitary certificate not required.
<i>Phytolacca</i>	No additional declarations required.
<i>Picea</i>	No additional declarations required.
<i>Pieris</i>	No additional declarations required.
<i>Pimpinella</i>	Phytosanitary certificate not required.
<i>Pinguicula</i>	Phytosanitary certificate not required.
<i>Piper</i>	No additional declarations required.
<i>Pisonia</i>	No additional declarations required.
<i>Pithecellobium</i>	No additional declarations required.
<i>Pittosporum</i>	No additional declarations required.
<i>Plagianthus</i>	No additional declarations required.
<i>Platycodon</i>	Phytosanitary certificate not required.
<i>Plectranthus</i>	No additional declarations required.
<i>Plumbago</i>	No additional declarations required.
<i>Plumeria</i>	No additional declarations required.
<i>Podocarpus</i>	No additional declarations required.
<i>Polemonium</i>	Phytosanitary certificate not required.
<i>Polyscias</i>	No additional declarations required.
<i>Pouteria</i>	No additional declarations required.
<i>Pothos</i>	Phytosanitary certificate not required.
<i>Pratia</i>	Phytosanitary certificate not required.
<i>Prieurella</i>	No additional declarations required.
<i>Primula</i>	Phytosanitary certificate not required.
<i>Prunella</i>	Phytosanitary certificate not required.
<i>Prumnopitys</i>	No additional declarations required.
<i>Pseudolachnostylis</i>	Phytosanitary certificate not required.
<i>Pseudolarix</i>	No additional declarations required.

<i>Pseudotsuga</i>	No additional declarations required.
<i>Psilotum</i>	Phytosanitary certificate not required.
<i>Psiloxylon</i>	No additional declarations required.
<i>Psophocarpus</i>	No additional declarations required.
<i>Psoralea</i>	Phytosanitary certificate not required.
<i>Pterocarpus</i>	No additional declarations required.
<i>Pterodiscus</i>	No additional declarations required.
<i>Puccinellia</i>	No additional declarations required.
<i>Puelia</i>	No additional declarations required.
<i>Pulicaria</i>	Phytosanitary certificate not required.
<i>Pulmonaria</i>	Phytosanitary certificate not required.
<i>Pultenaea</i>	No additional declarations required.
<i>Punica</i>	No additional declarations required.
<i>Pyrostegia</i>	No additional declarations required.
<i>Radermachera</i>	No additional declarations required.
<i>Raphionacme</i>	No additional declarations required.
<i>Ravenala</i>	No additional declarations required.
<i>Rehmannia</i>	No additional declarations required.
<i>Reseda</i>	Phytosanitary certificate not required.
<i>Restio</i>	No additional declarations required.
<i>Reyesia</i>	Phytosanitary certificate not required.
<i>Rhamnus</i>	No additional declarations required.
<i>Rhodochiton</i>	No additional declarations required.
<i>Rhododendron</i>	No additional declarations required.
<i>Rhodohypoxis</i>	Phytosanitary certificate not required.
<i>Rhodosphaera</i>	No additional declarations required.
<i>Rhoeo</i>	Phytosanitary certificate not required.
<i>Rhytidocaulon</i>	No additional declarations required.
<i>Rogersia</i>	Phytosanitary certificate not required.
<i>Romneya</i>	Phytosanitary certificate not required.
<i>Roscoea</i>	No additional declarations required.
<i>Rosmarinus</i>	No additional declarations required.

<i>Rothmannia</i>	No additional declarations required.
<i>Rubia</i>	Phytosanitary certificate not required.
<i>Rudbeckia</i>	No additional declarations required.
<i>Ruellia</i>	Phytosanitary certificate not required.
<i>Ryania</i>	Phytosanitary certificate not required.
<i>Sagina</i>	Phytosanitary certificate not required.
<i>Salix</i>	No additional declarations required.
<i>Salpiglossus</i>	Phytosanitary certificate not required.
<i>Salvia</i>	Phytosanitary certificate not required.
<i>Samanea</i>	No additional declarations required.
<i>Sandoricum</i>	No additional declarations required.
<i>Santalum</i>	No additional declarations required.
<i>Sanvitalia</i>	Phytosanitary certificate not required.
<i>Saponaria</i>	Phytosanitary certificate not required.
<i>Sapium</i>	No additional declarations required.
<i>Saraca</i>	No additional declarations required.
<i>Sarcocephalus</i>	No additional declarations required.
<i>Sarcococca</i>	No additional declarations required.
<i>Sarracenia</i>	Phytosanitary certificate not required.
<i>Satureja</i>	Phytosanitary certificate not required.
<i>Saxifraga</i>	Phytosanitary certificate not required.
<i>Scabiosa</i>	Phytosanitary certificate not required.
<i>Scaevola</i>	No additional declarations required.
<i>Schefflera</i>	No additional declarations required.
<i>Schinus</i>	No additional declarations required.
<i>Schizaea</i>	Phytosanitary certificate not required.
<i>Schizanthus</i>	Phytosanitary certificate not required.
<i>Schizolobium</i>	No additional declarations required.
<i>Scindapsus</i>	Phytosanitary certificate not required.
<i>Scorzomera</i>	Phytosanitary certificate not required.
<i>Schoenocaulon</i>	No additional declarations required.
<i>Schotia</i>	No additional declarations required.

<i>Securidaca</i>	No additional declarations required.
<i>Sempervivum</i>	Phytosanitary certificate not required.
<i>Sesamum</i>	No additional declarations required.
<i>Shasta</i>	Phytosanitary certificate not required.
<i>Shepherdia</i>	No additional declarations required.
<i>Sicana</i>	No additional declarations required.
<i>Sidalcea</i>	Phytosanitary certificate not required.
<i>Silene</i>	Phytosanitary certificate not required.
<i>Simmondsia</i>	No additional declarations required.
<i>Sinapis</i>	No additional declarations required.
<i>Soleirolia</i>	No additional declarations required.
<i>Solidago</i>	Phytosanitary certificate not required.
<i>Solidaster</i>	Phytosanitary certificate not required.
<i>Solenostemon</i>	Phytosanitary certificate not required.
<i>Sollya</i>	No additional declarations required.
<i>Sophora</i>	No additional declarations required.
<i>Sparmannia</i>	No additional declarations required.
<i>Spartium</i>	No additional declarations required.
<i>Spathiphyllum</i>	No additional declarations required.
<i>Spiraea</i>	No additional declarations required.
<i>Spondias</i>	No additional declarations required.
<i>Spyridium</i>	No additional declarations required.
<i>Stachys</i>	Phytosanitary certificate not required.
<i>Stenocarpus</i>	No additional declarations required.
<i>Stephanotis</i>	No additional declarations required.
<i>Stereospermum</i>	No additional declarations required.
<i>Stevia</i>	Phytosanitary certificate not required.
<i>Stizolobium</i>	Phytosanitary certificate not required.
<i>Stokesia</i>	Phytosanitary certificate not required.
<i>Streptocarpus</i>	Phytosanitary certificate not required.
<i>Strychnos</i>	No additional declarations required.
<i>Stylosanthes</i>	No additional declarations required.

<i>Styphelia</i>	No additional declarations required.
<i>Swainsona</i>	No additional declarations required.
<i>Swartzia</i>	No additional declarations required.
<i>Swientenia</i>	No additional declarations required.
<i>Symphoricarpos</i>	No additional declarations required.
<i>Symphytum</i>	Phytosanitary certificate not required.
<i>Syngonium</i>	Phytosanitary certificate not required.
<i>Synsepalum</i>	No additional declarations required.
<i>Syzygium</i>	No additional declarations required.
<i>Tabebuia</i>	No additional declarations required.
<i>Tabernaemontana</i>	No additional declarations required.
<i>Tacca</i>	Phytosanitary certificate not required.
<i>Talinum</i>	Phytosanitary certificate not required.
<i>Tamarindus</i>	No additional declarations required.
<i>Tamarix</i>	No additional declarations required.
<i>Tanacetum</i>	Phytosanitary certificate not required.
Taxodiaceae	No additional declarations required.
<i>Telopea</i>	No additional declarations required.
<i>Templetonia</i>	No additional declarations required.
<i>Tephrosia</i>	No additional declarations required.
<i>Tetragonia</i>	Phytosanitary certificate not required.
<i>Tetraclinis</i>	No additional declarations required.
<i>Tetrataxis</i>	No additional declarations required.
<i>Teucrium</i>	Phytosanitary certificate not required.
<i>Thalictrum</i>	Phytosanitary certificate not required.
<i>Thermopsis</i>	Phytosanitary certificate not required.
<i>Thespesia</i>	No additional declarations required.
<i>Thunbergia</i>	No additional declarations required.
<i>Thymophylla</i>	Phytosanitary certificate not required.
<i>Thymus</i>	Phytosanitary certificate not required.
<i>Tiarella</i>	Phytosanitary certificate not required.
<i>Tibouchina</i>	Phytosanitary certificate not required.

<i>Tilia</i>	No additional declarations required.
<i>Tmesipterus</i>	No additional declarations required.
<i>Torenia</i>	Phytosanitary certificate not required.
<i>Trachelium</i>	Phytosanitary certificate not required.
<i>Trachymene</i>	Phytosanitary certificate not required.
<i>Tradescantia</i>	Phytosanitary certificate not required.
<i>Tragopogon</i>	Phytosanitary certificate not required.
<i>Triaspis</i>	No additional declarations required.
<i>Trigonella</i>	Phytosanitary certificate not required.
<i>Tristania</i>	No additional declarations required.
<i>Tritonia</i>	Phytosanitary certificate not required.
<i>Tropaeolum</i>	Phytosanitary certificate not required.
<i>Tsuga</i>	No additional declarations required.
<i>Turraea</i>	No additional declarations required.
<i>Umbellularia</i>	No additional declarations required.
<i>Uncarina</i>	No additional declarations required.
<i>Ursinia</i>	Phytosanitary certificate not required.
<i>Utricularia</i>	Phytosanitary certificate not required.
<i>Valeriana</i>	No additional declarations required.
<i>Valerianella</i>	Phytosanitary certificate not required.
<i>Vangueria</i>	No additional declarations required.
<i>Verbascum</i>	No additional declarations required.
<i>Verbena</i>	Phytosanitary certificate not required.
<i>Vernonia</i>	No additional declarations required.
<i>Veronica</i>	Phytosanitary certificate not required.
<i>Verticordia</i>	No additional declarations required.
<i>Viburnum</i>	No additional declarations required.
<i>Victoria</i>	No additional declarations required.
<i>Villarisa</i>	Phytosanitary certificate not required.
<i>Vinca</i>	Phytosanitary certificate not required.
<i>Viscaria</i>	Phytosanitary certificate not required.
<i>Vitex</i>	No additional declarations required.

<i>Voandzeia</i>	Phytosanitary certificate not required.
<i>Wahlenbergia</i>	Phytosanitary certificate not required.
<i>Wasabia</i>	Phytosanitary certificate not required.
<i>Warburgia</i>	No additional declarations required.
<i>Weinmannia</i>	No additional declarations required.
<i>Welwitschia</i>	No additional declarations required.
<i>Wisteria</i>	No additional declarations required.
<i>Witheringia</i>	No additional declarations required.
<i>Xanthisma</i>	Phytosanitary certificate not required.
<i>Xeranthemum</i>	Phytosanitary certificate not required.
<i>Zinnia</i>	Phytosanitary certificate not required.

4.6 Herbs, Spices, Medicinal Plants, & Plants for Animal & Human Consumption (Including Seed for Consumption)

4.6.1 Herbs, Spices, Medicinal Plants, & Plants for Animal & Human Consumption (Including Seed for Consumption)

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Herbs, Spices, Medicinal Plants, and Plants for Animal & Human Consumption (Including Seed for Consumption) require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Acanthus mollis</i>	No additional declarations required.
<i>Achillea</i>	No additional declarations required.
<i>Aconitum</i>	No additional declarations required.
<i>Actaea spicata</i>	No additional declarations required.
<i>Agar</i>	No additional declarations required.
<i>Agastache foeniculum</i>	No additional declarations required.
<i>Agrimonia eupatoria</i>	No additional declarations required.
<i>Ajuga</i>	No additional declarations required.
<i>Alcea rosea</i>	No additional declarations required.
<i>Alchemilla vulgaris</i>	No additional declarations required.

<i>Allium</i> (dried, flaked, powdered)	No additional declarations required.
<i>Aloe</i>	No additional declarations required.
<i>Althaea</i>	No additional declarations required.
<i>Anacardium</i> (nuts only)	No additional declarations required.
<i>Anagallis arvensis</i>	No additional declarations required.
<i>Anchusa</i>	No additional declarations required.
<i>Androstephium violaceum</i>	No additional declarations required.
<i>Anemone</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Anethum</i>	No additional declarations required.
<i>Angelica</i>	No additional declarations required.
<i>Antennaria dioica</i>	No additional declarations required.
<i>Anthemis tinctoria</i>	No additional declarations required.
<i>Anthriscus</i>	No additional declarations required.
<i>Anthyllis</i>	No additional declarations required.
<i>Apium graveolens</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Aquilegia vulgaris</i>	No additional declarations required.
<i>Arabis</i>	No additional declarations required.
<i>Aralia</i>	No additional declarations required.
<i>Arctotis acaulis</i>	No additional declarations required.
<i>Areca catechu</i> (nuts only)	No additional declarations required.
<i>Armeria</i>	No additional declarations required.
<i>Arnica montana</i>	No additional declarations required.
<i>Asarum canadense</i>	No additional declarations required.
<i>Asclepias tuberosa</i>	No additional declarations required.
<i>Asperula</i>	No additional declarations required.
<i>Astragalus</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Atropa</i>	No additional declarations required.
<i>Baptisia</i>	No additional declarations required.
<i>Barbarea verna</i>	No additional declarations required.
<i>Bellis perennis</i>	No additional declarations required.
<i>Borago officinalis</i>	No additional declarations required.

<i>Brassica</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Bryonia dioica</i>	No additional declarations required.
<i>Cajanus</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Calendula</i>	No additional declarations required.
<i>Campanula</i>	No additional declarations required.
<i>Canavalia</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Capparis spinosa</i>	No additional declarations required.
<i>Carthamus tinctorius</i>	No additional declarations required.
<i>Carum carvi</i>	No additional declarations required.
<i>Carya</i> (nuts only)	No additional declarations required.
<i>Castanea</i>	No additional declarations required.
<i>Catharanus roseus</i>	No additional declarations required.
<i>Ceanothus americanus</i>	No additional declarations required.
<i>Cedronella tryphylla</i>	No additional declarations required.
<i>Centaurea cyanus</i>	No additional declarations required.
<i>Centaurium erythraea</i>	No additional declarations required.
<i>Ceratonia siliqua</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cheiranthus cheiri</i>	No additional declarations required.
<i>Chelidonium</i>	No additional declarations required.
<i>Chenopodium</i>	No additional declarations required.
<i>Chrysanthemum</i>	No additional declarations required.
<i>Cicer</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cichorium intybus</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Cimicifuga racemosa</i>	No additional declarations required.

<i>Cinnamomum</i>	No additional declarations required.
<i>Cnicus benedictus</i>	No additional declarations required.
<i>Cocos nucifera</i> (nuts only)	No additional declarations required.
<i>Coffea</i> (Roasted beans only)	No additional declarations required.
<i>Colchicum autumnale</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Collinsonia canadensis</i>	No additional declarations required.
<i>Colutea</i>	No additional declarations required.
<i>Conium maculatum</i>	No additional declarations required.
<i>Convallaria</i>	No additional declarations required.
<i>Coreopsis tinctoria</i>	No additional declarations required.
<i>Coriandrum sativum</i>	No additional declarations required.
<i>Cornus florida</i>	No additional declarations required.
<i>Corylus</i> (nuts only)	No additional declarations required.
<i>Crithmum maritimum</i>	No additional declarations required.
<i>Crocus sativus</i>	No additional declarations required.
<i>Croton</i>	No additional declarations required.
<i>Cryptotaenia japonica</i>	No additional declarations required.
Cucurbitaceae (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cuminum cyminum</i>	No additional declarations required.
<i>Cupressus</i>	No additional declarations required.
<i>Curcuma longa</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Cyamopsis</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cynara</i>	No additional declarations required.
<i>Danae racemosa</i>	No additional declarations required.
<i>Delphinium grandiflorum</i>	No additional declarations required.
<i>Dianthus</i>	No additional declarations required.
<i>Dictamnus albus</i>	No additional declarations required.
<i>Digitalis</i>	No additional declarations required.

<i>Dipsacus</i> (flower head)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Dolichos</i>	No additional declarations required.
<i>Dracocephalum moldavica</i>	No additional declarations required.
<i>Elettaria cardemomum</i>	No additional declarations required.
<i>Epilobium</i>	No additional declarations required.
<i>Erica</i>	No additional declarations required.
<i>Erigeron</i>	No additional declarations required.
<i>Eugenia caryophyllata</i>	No additional declarations required.
<i>Euphorbia</i>	No additional declarations required.
<i>Fagopyrum</i>	No additional declarations required.
<i>Filipendula</i>	No additional declarations required.
<i>Foeniculum vulgare</i>	No additional declarations required.
<i>Galega officinale</i>	No additional declarations required.
<i>Galium</i>	No additional declarations required.
<i>Genista tinctoria</i>	No additional declarations required.
<i>Gentiana</i>	No additional declarations required.
<i>Geranium</i>	No additional declarations required.
<i>Glycines</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Glycyrrhiza glabra</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Gnaphalium</i>	No additional declarations required.
<i>Gossypium</i>	No additional declarations required. T2 = consignment was treated with a wide spectrum insecticide or fumigant. D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.
<i>Gratiola officinalis</i>	No additional declarations required.
<i>Grindelia robusta</i>	No additional declarations required.
<i>Guizotia abyssinica</i>	No additional declarations required.
<i>Hamamelis virginiana</i>	No additional declarations required.

<i>Helianthemum</i>	No additional declarations required.
<i>Heliotropium</i>	No additional declarations required.
<i>Helleborus niger</i>	No additional declarations required.
<i>Herniaria</i>	No additional declarations required.
<i>Heuchera</i>	No additional declarations required.
<i>Hibiscus sabdariffa</i>	No additional declarations required.
<i>Hieracium</i>	No additional declarations required.
<i>Hordeum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Humulus</i>	No additional declarations required.
<i>Hydrocotyle asiatica</i>	No additional declarations required.
<i>Hyoscyamus niger</i>	No additional declarations required.
<i>Hypericum</i>	No additional declarations required.
<i>Hyssopus officinalis</i>	No additional declarations required.
<i>Iberis umbellata</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Ilex paraguariensis</i>	No additional declarations required.
<i>Illicium verum</i>	No additional declarations required.
<i>Indigofera</i>	No additional declarations required.
<i>Inula helenium</i>	No additional declarations required.
<i>Iris</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Isatis tinctoria</i>	No additional declarations required.
<i>Jasminum</i>	No additional declarations required.
<i>Juglans</i> (nuts only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Juniperus communis</i>	No additional declarations required.
<i>Lamium</i>	No additional declarations required.
<i>Laurus</i>	No additional declarations required.
<i>Lavandula</i>	No additional declarations required.
<i>Lawsona inermis</i>	No additional declarations required.

<i>Lens</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Leontopodium alpinum</i>	No additional declarations required.
<i>Leonurus</i>	No additional declarations required.
<i>Lepidium sativum</i>	No additional declarations required.
<i>Levisticum officinale</i>	No additional declarations required.
<i>Liatris</i>	No additional declarations required.
<i>Ligustrum</i>	No additional declarations required.
<i>Linaria</i>	No additional declarations required.
<i>Linum usitatissimum</i>	No additional declarations required.
<i>Lobelia</i>	No additional declarations required.
<i>Lophocarpus</i>	No additional declarations required.
<i>Lotus</i>	No additional declarations required.
<i>Lupinus</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Lychnis chalcedonica</i>	No additional declarations required.
<i>Lycium chinense</i>	No additional declarations required.
<i>Lycopus europaeus</i>	No additional declarations required.
<i>Lysimachia nummularia</i>	No additional declarations required.
<i>Lythrum salicaria</i>	No additional declarations required.
<i>Macadamia</i> (nuts only)	No additional declarations required.
<i>Maclura pomifera</i>	No additional declarations required.
<i>Mangifera</i> (seed only)	No additional declarations required.
<i>Malva</i>	No additional declarations required.
<i>Manihot esculenta</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Maranta arundinacea</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Marrubium vulgare</i>	No additional declarations required.
<i>Matricaria chamomilla</i>	No additional declarations required.
<i>Melilotus officinalis</i>	No additional declarations required.
<i>Melissa officinalis</i>	No additional declarations required.
<i>Mentha</i>	No additional declarations required.

<i>Metroxylon rumphii</i>	No additional declarations required.
<i>Mirabilis</i>	No additional declarations required.
<i>Molucella laevis</i>	No additional declarations required.
<i>Monarda didyma</i>	No additional declarations required.
<i>Morus nigra</i> (dried)	No additional declarations required.
<i>Myrica pennsylvanica</i>	No additional declarations required.
<i>Myristica fragrans</i>	No additional declarations required.
<i>Myrrhis odorata</i>	No additional declarations required.
<i>Myrtus communis</i>	No additional declarations required.
<i>Nasturtium officinale</i>	No additional declarations required.
<i>Nepenthes</i>	No additional declarations required.
<i>Nepeta</i>	No additional declarations required.
<i>Nigella damascena</i>	No additional declarations required.
<i>Ocimum</i>	No additional declarations required.
<i>Ononis</i>	No additional declarations required.
<i>Origanum</i>	No additional declarations required.
<i>Oryza</i> (grain only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Panax</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Papaver somniferum</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with MBr gas. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Paris quadrifolia</i>	No additional declarations required.
<i>Pelargonium</i>	No additional declarations required.
<i>Perilla frutescens</i>	No additional declarations required.
<i>Petroselinum crispum</i>	No additional declarations required.
<i>Peucedanum</i>	No additional declarations required.
<i>Phalaris canariensis</i>	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Phaseolus</i>	No additional declarations required.
<i>Piper</i>	No additional declarations required.

<i>Pisum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Ricinus</i>	No additional declarations required.
<i>Saxifraga</i>	No additional declarations required.
<i>Secale</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Sedum</i>	No additional declarations required.
<i>Sempervivum</i>	No additional declarations required.
<i>Sesamum</i>	No additional declarations required.
<i>Silybum marianum</i>	No additional declarations required.
<i>Smilax</i>	No additional declarations required.
<i>Solidago virgaurea</i>	No additional declarations required.
<i>Sorghum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Stachys</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Symphytum</i>	No additional declarations required.
<i>Tagetes</i>	No additional declarations required.
<i>Taraxacum officinale</i>	No additional declarations required.
<i>Teucrium</i>	No additional declarations required.
<i>Tetragonia tetragonioides</i>	No additional declarations required.
<i>Theobroma</i>	No additional declarations required.
<i>Thymus</i>	No additional declarations required.
<i>Tilia cordata</i>	No additional declarations required.
<i>Trigonella foenum-graecum</i>	No additional declarations required.
<i>Tropaeolum</i>	No additional declarations required.
<i>Tussilago farfara</i>	No additional declarations required.
<i>Urginea maritima</i>	No additional declarations required.
<i>Urtica</i>	No additional declarations required.
<i>Valeriana</i>	No additional declarations required.
<i>Valerianella locusta</i>	No additional declarations required.

<i>Vanilla</i>	No additional declarations required.
<i>Verbascum</i>	No additional declarations required.
<i>Verbena</i>	No additional declarations required.
<i>Vernonia</i>	No additional declarations required.
<i>Veronica</i>	No additional declarations required.
<i>Viburnum opulus</i>	No additional declarations required.
<i>Vigna</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Vinca</i>	No additional declarations required.
<i>Viola</i>	No additional declarations required.
<i>Voandzeia</i>	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Yucca glauca</i>	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Zea</i>	AD2:- New Zealand is free from <i>Cephalosporium maydis</i> , and <i>Peronosclerospora maydis</i> . AD3:- The area of production is free from <i>Erwinia stewartii</i> . AD4:- The consignment is free from <i>Prostephanus truncatus</i> . (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)