



**Review of Yersiniosis Notifications in New Zealand
2002-2006**

Prepared as part of a New Zealand Food Safety Authority
contract for scientific services

by

Ruth Pirie
Jonathan Williman
Carolyn Nicol
Kerry Sexton

April 2008

Client Report
FW07111

**Review of Yersiniosis Notifications in New Zealand
2002-2006**

Bruce Adlam

Population and Environmental Health Programme Leader

Ruth Pirie

Project Leader

Bruce Adlam

Peer Reviewer

DISCLAIMER

This report or document (“the Report”) is given by the Institute of Environmental Science and Research Limited (“ESR”) solely for the benefit of the New Zealand Food Safety Authority (“NZFSA”), Public Health Services Providers and other Third Party Beneficiaries as defined in the Contract between ESR and the NZFSA, and is strictly subject to the conditions laid out in that Contract.

Neither ESR nor any of its employees makes any warranty, express or implied, or assumes any legal liability or responsibility for use of the Report or its contents by any other person or organisation.

ACKNOWLEDGEMENTS

This report could not have been completed without the assistance of the staff in public health offices that enter information disease notifications into EpiSurv and those who completed and returned the questionnaires.

The authors would also like to thank Dinusha Bandara at ESR for her help with the statistical tests and Dr Donald Campbell at the New Zealand Food Safety Authority for his guidance and helpful review during the project.

CONTENTS

| | | |
|----------|---|-----------|
| 1 | SUMMARY | 6 |
| 2 | INTRODUCTION | 7 |
| 3 | METHODS | 9 |
| 3.1 | Datasets used in the analyses | 9 |
| 3.2 | Medical Officers of Health survey | 10 |
| 3.3 | Case definition for yersiniosis notification | 10 |
| 3.4 | Pathogenicity of <i>Yersinia</i> spp. | 10 |
| 3.5 | Statistical tests..... | 11 |
| | RESULTS | 12 |
| 3.6 | EpiSurv notification data trends..... | 12 |
| 3.7 | Source of notifications | 14 |
| 3.8 | Cases meeting yersiniosis case definition..... | 14 |
| 3.9 | Enteric Reference Laboratory (ERL) <i>Yersinia</i> biotyping data..... | 16 |
| 3.10 | Hospitalisations..... | 23 |
| 3.11 | <i>Yersinia</i> testing methodologies used by clinical laboratories in New Zealand..... | 25 |
| 3.12 | Medical Officers of Health Survey | 26 |
| 3.12.1 | Source of notification..... | 26 |
| 3.12.2 | Investigation of cases | 26 |
| 3.12.3 | Local trends in yersiniosis notifications | 27 |
| 3.12.4 | <i>Yersinia</i> strain and biotyping information | 27 |
| 3.12.5 | De-notification of yersiniosis cases | 27 |
| 3.12.6 | Other Comments | 27 |
| 4 | DISCUSSION | 28 |
| 5 | RECOMMENDATIONS | 30 |
| 6 | REFERENCES | 31 |
| | APPENDIX A – MEDICAL OFFICERS OF HEALTH YERSINIOSIS SURVEY RESPONDENTS | 32 |
| | APPENDIX B – YERSINIOSIS QUESTIONNAIRE | 33 |

LIST OF TABLES

| | |
|--|----|
| Table 1. DALYs for infectious intestinal diseases in New Zealand | 7 |
| Table 2. Annual incidence of disease caused by foodborne bacterial agents in OECD countries | 7 |
| Table 3. Biotype, O serotype and pYV carriage of <i>Y. enterocolitica</i> | 11 |
| Table 4. Categories used to grouping <i>Yersinia</i> strains by pathogenicity | 11 |
| Table 5. Number of cases and rates per 100,000 of yersiniosis notifications by DHB | 13 |
| Table 6. Number and rates per 100,000 of notifiable yersiniosis cases by age group | 14 |
| Table 7. Number and percentage of notifiable yersiniosis cases by reporting source | 14 |
| Table 8. Number and percentage of notifiable yersiniosis cases by case status | 15 |
| Table 9. Number and percentage of notifiable confirmed yersiniosis cases reported meeting clinical criteria of case definition | 15 |
| Table 10. Number of isolates typed by ERL by strain of <i>Yersinia</i> as a percentage of all <i>Yersinia</i> isolates typed that year. | 16 |
| Table 11. Number and percentage of isolates typed by ERL by biotype of <i>Y. enterocolitica</i> . 16 | |
| Table 12. Number and percentage of isolates typed by ERL and matched to an EpiSurv case | 18 |
| Table 13. Number of isolates typed by ERL and matched to an EpiSurv case as a percentage of all isolates typed that year, by DHB | 18 |
| Table 14. Number and percentage of <i>Yersinia</i> isolates typed by ERL and matched to an EpiSurv case that meet the current laboratory criteria for yersiniosis notification (<i>Y. enterocolitica</i> or <i>Y. pseudotuberculosis</i>) by DHB | 19 |
| Table 15. Summary of notifications and laboratory biotyping results, 2002 to 2006 | 20 |
| Table 16. Summary of notifications and laboratory biotyping results by DHB for the period 2002 to 2006 | 21 |
| Table 17. Number of yersiniosis notifications reported on EpiSurv, <i>Yersinia</i> isolates typed by the ERL and EpiSurv records updated with <i>Yersinia</i> strain by DHB and year, 2002 to 2006. 22 | |
| Table 18. Number and percentage of notified yersiniosis cases admitted to a hospital reported on EpiSurv | 23 |
| Table 19. Number of hospitalised yersiniosis cases (ICD10 A046) by diagnosis status, as reported by NZHIS | 23 |
| Table 20. Number and percentage of hospitalised yersiniosis reported by NZHIS that could be matched to cases on EpiSurv and isolates typed by ERL | 23 |
| Table 21. <i>Yersinia</i> strain and biotype (from ERL) for NZHIS reported yersiniosis cases by diagnostic status | 24 |
| Table 22. Number of hospitalised yersiniosis cases reported on EpiSurv compared to NZHIS reported cases. | 25 |

LIST OF FIGURES

| | |
|---|----|
| Figure 1. Yersiniosis notifications by year, 2002-2006..... | 12 |
| Figure 2. Average number of cases reported per month, 2002 to 2006..... | 12 |
| Figure 3. Graph of notifications by pathogenic group of <i>Yersinia</i> by year, 2002-2006..... | 17 |
| Figure 4. Graph of <i>Yersinia</i> isolates by year, quarter and pathogenic grouping, 2002-2006.. | 17 |

1 SUMMARY

In the five year period (2002 to 2006) reviewed, the annual notification rate for yersiniosis in New Zealand increased over the four years 2003 to 2006 but remained lower than it was in 2002. A number of District Health Boards (DHBs) (West Coast, South Canterbury, Capital and Coast) had consistently higher notification rates than the overall New Zealand notification rate. Children aged less than 5 years experienced the highest rates of yersiniosis and there was a small increase in the notification rate for the 60 years and over age group.

Not all *Yersinia* spp. are pathogenic and the biotyping data is a very useful tool to investigate notification practices. There was a large variation in the practices of laboratories between DHBs in referring isolates to ESR's Enteric Reference Laboratory for characterisation e.g. laboratories in some DHBs sent in more than 50% of their isolates while in other DHBs no isolates were submitted in the five year period. A feature of the biotyping data in more recent years was the increasing percentage of non pathogenic isolates being typed. Public Health Services (PHSs) requested guidance in the interpretation and use of *Yersinia* biotyping results.

An increasing number of hospitalisations for yersiniosis were reported over the five years reviewed with approximately half of these being admissions for yersiniosis as a primary diagnosis. The numbers were very small but the combining of the yersiniosis hospitalisations data with the *Yersinia* strain and biotyping information showed that an increasing number of the hospitalisations were for non pathogenic strains of *Yersinia*.

PHSs reported varying practices in the investigation of yersiniosis cases with most PHSs investigating yersiniosis cases to some extent. Biotyping results did not appear to be readily available to PHSs and often did not make their way into EpiSurv. The new requirement from 18 December 2007 for laboratories to directly report notifiable diseases to Medical Officers of Health should improve this situation.

2 INTRODUCTION

Yersiniosis is the third most commonly reported potential food-borne disease notified in New Zealand. These numbers have had a major influence on risk ranking exercises carried out by the New Zealand Food Safety Authority (NZFSA) with yersiniosis being estimated to contribute a higher number of disability adjusted life years (DALYs) to the New Zealand burden of food-borne disease than *E. coli* O157 infection (Table 1).

Table 1. DALYs for infectious intestinal diseases in New Zealand

| Disease | DALYs |
|------------------------|-------|
| Campylobacteriosis | 1554 |
| Norovirus infection | 536 |
| Listeriosis, perinatal | 229 |
| Salmonellosis | 186 |
| Yersiniosis | 93 |
| STEC infection | 91 |
| Listeriosis, acquired | 26 |

Source: [1]

New Zealand's notification rate for yersiniosis in 2006 (11.8 cases per 100,000 population) was high compared to other OECD countries though lower than Finland.

Table 2. Annual incidence of disease caused by foodborne bacterial agents in OECD countries

| Country | Year | Cases | Incidence (per 100 000 population) |
|----------------|------|----------------|---------------------------------------|
| Australia | 2000 | 73 | 0.6 |
| Austria | 1998 | 94 | 1.2 |
| Belgium | 2000 | 507 | 5 |
| Denmark | 2001 | 286 | 5.3 |
| Finland | 2001 | 728 | 14 |
| Greece | 1998 | 10 | 0.1 |
| Japan | 2001 | 4 | <0.01 |
| Norway | 2001 | 123 | 2.8 |
| Spain | 1998 | 425 | 1.1 |
| Sweden | 2001 | 579 | 6.5 |
| Switzerland | 1998 | 51 | 0.7 |
| United Kingdom | 2000 | 27 | 0.05 |
| United States | 1999 | - ^a | 0.4 |

^a Data not provided

Source : [2]

In New Zealand there was an apparent surge in yersiniosis cases reported in late 2006, especially in the population served by Christchurch Hospital. Anecdotal evidence suggests that there may have been differing clinical practices (e.g. infection screening versus clinical diagnosis), serology reporting and use of reference services occurring throughout the country leading to differential notification.

In light of both the disease burden and its consequent implications for trade, ESR was contracted by NZFSA to ascertain the validity of the yersiniosis notification data.

The objectives of this project were

- To validate the human yersiniosis disease notification data held in the national notifiable disease database, EpiSurv.
- To validate an agreed sample (time and geography) of yersiniosis notifications utilising notification, clinical, clinical laboratory and reference laboratory information
- To produce a report with recommendations if appropriate based on the findings.

3 METHODS

Three sources of data were used to carry out this review. Yersiniosis notifications held in EpiSurv were used as the primary dataset with supplementary information being obtained by matching notifications to data held by ESR's Enteric Reference Laboratory (ERL) and New Zealand Health Information Service (NZHIS) hospitalisation data.

3.1 Datasets used in the analyses

a) EpiSurv notifiable disease data

Yersiniosis notification data for all of New Zealand was extracted from the EpiSurv database on the 18 October 2007 for the period 1 January 2002 to 31 December 2006, with the following variables:

- EpiSurv identification number
- Case status
- Fits clinical description
- Meets laboratory criteria
- Reporting source (GP, Hospital, Laboratory, Self-reported, Outbreak, Other)
- Was the case hospitalised
- Hospitalisation date
- District Health Board
- Age
- Sex
- Ethnicity (Prioritised)
- Reporting date
- Species and strain information from laboratory

Rates were calculated using Statistics New Zealand mid year population estimates.

b) ERL

Yersinia isolates from cases may be submitted by clinical laboratories for ERL to determine strain and biotype information. This information, stored by ESR in ESRLab, was matched to notifications reported on EpiSurv using case names, sex, date of birth, and report date of disease.

c) New Zealand Health Information Service

Hospitalisation data for cases of "Enteritis due to *Yersinia enterocolitica*" (ICD-10 code A04.6) for all of New Zealand were obtained from NZHIS for the period 1 January 2002 to 31 December 2006 with the following variables:

- Date of admission
- Age
- Gender
- Ethnicity grouping
- District Health Board
- Diagnosis type

Cases were matched to notifications on EpiSurv using age, ethnicity, District Health Board and report date. Cases were also matched to isolates received by ERL either indirectly through their

matched EpiSurv record, or directly using sex, age, date of hospitalisation/date isolate received, DHB and client laboratory.

3.2 Medical Officers of Health survey

A survey of Medical Officers of Health (MOsH) was undertaken to identify differences between PHSs in the handling of yersiniosis cases, in particular what practices were in use to determine whether yersiniosis notifications met the current case definition.

The survey was e-mailed to each public health office with a Medical Officer of Health. Every public health office returned a completed questionnaire giving a 100 % response rate (17 out of 17 public health offices). A list of the respondents can be found in Appendix A and a copy of the questionnaire can be found in Appendix B.

3.3 Case definition for yersiniosis notification

Cases were classified by PHSs as confirmed or probable according to the case definition for yersiniosis in the CDC manual.[3] The case definition included a clinical and laboratory component as shown in below.

Clinical description

An acute illness with diarrhoea, fever and abdominal pain. Mesenteric adenitis may occur and complications include arthritis and systemic infection.

Laboratory test for diagnosis

Isolation of *Yersinia enterocolitica* or *Y. pseudotuberculosis* from blood or faeces

OR

Detection of circulating antigen by ELISA or agglutination test.

Case classification

Probable: A clinically compatible illness that is epidemiologically linked to a confirmed case.

Confirmed: A clinically compatible illness that is laboratory confirmed.

3.4 Pathogenicity of *Yersinia* spp.

Of the 11 species within the *Yersinia* genus only three, *Y. pestis*, *Y. pseudotuberculosis* and *Y. enterocolitica*, are regarded as pathogenic to humans. [4]

Y. pestis is the causative agent of bubonic and pneumonic plague, *Y. pseudotuberculosis* is a rodent pathogen which occasionally causes mesenteric lymphadenitis, septicaemia and immune-mediated diseases in humans and *Y. enterocolitica* is a versatile intestinal pathogen which is the most prevalent *Yersinia* species amongst humans.

Of the three pathogenic species of *Yersinia*, *Y. enterocolitica* is the most heterogeneous and can be divided into approximately 30 distinct serotypes (based on antigenic variation in cell wall lipopolysaccharide) and six biotypes (based on variations in biochemical behaviour). Table 3 shows the relationship between biotype, O serotype and pYV (*Yersinia* virulence plasmid) carriage of *Y. enterocolitica*.

Table 3. Biotype, O serotype and pYV carriage of *Y. enterocolitica*

| Biotype | Serotype(s) |
|----------------|--|
| 1A | O:4; O:5; O:6,30; O6,31; O:7,8; O:7,13; O:10; O:14; O:16; O:21; O:22; O:25; O:37; O:41,42; O:46; O:47; O:57; NT ^a |
| 1B | O:4,32 ^b ; O:8 ^b ; O:13 ^a ,13b ^b ; O:16; O:18 ^b ; O:20 ^b ; O:21 ^b ; O:25; O:41,42; NT |
| 2 | O:5,27 ^b ; O:9 ^b ; O:27 |
| 3 | O:1,2,3 ^b ; O:3 ^b ; O:5,27 ^b |
| 4 | O:3 ^b |
| 5 | O:2,3 ^b |

^a NT, not typable.

^b Serotypes which include strains that carry pYV.

Source: [5]

Y. enterocolitica strains of biotype 1A lack the known virulence determinants of strains in other categories, including the pYV *Yersinia* virulence plasmid, and several chromosomal markers of pathogenicity. For this reason, and also because *Y. enterocolitica* strains of biotype 1A are frequently isolated from the environment or asymptomatic individuals, these bacteria are often assumed to be avirulent.

However there is a considerable body of clinical, epidemiological and experimental evidence to indicate that at least some strains of *Y. enterocolitica* biotype 1A are able to cause gastrointestinal symptoms which resemble those caused by pYV-bearing strains.[5]

The categories shown in Table 4 are used in this report to assist in the analysis of trends in the reporting of yersiniosis and distinguish between the pathogenicity of the different strains of *Yersinia*.

Table 4. Categories used to group *Yersinia* strains by pathogenicity

| Category | Description |
|-----------------|--|
| Non pathogenic | All <i>Yersinia</i> species excluding <i>Y. enterocolitica</i> , <i>Y. pseudotuberculosis</i> and <i>Y. pestis</i> |
| Biotype 1A | <i>Y. enterocolitica</i> biotype 1A which are potentially but often not pathogenic |
| Pathogenic | <i>Y. enterocolitica</i> biotypes 1B, 2, 3, 4 and 5 <i>Y. pseudotuberculosis</i> and <i>Y. pestis</i> ^a |
| Unknown | No strain or biotyping information available |

^a No notifications of *Y. pestis* were reported in EpiSurv during the study period.

3.5 Statistical tests


The Mantel-Haenszel chi-square test was used to determine statistical significance. P-values less than 0.05 are considered to be significant at the 95% level of confidence. Fisher's Exact Test was also considered when 25% of the cells had expected counts less than 5.

RESULTS

3.6 EpiSurv notification data trends


The number of yersiniosis notifications reported in New Zealand decreased from 476 in 2002 to 407 in 2005 before increasing to 487 in 2006 (Figure 1). The national population rate of yersiniosis varied between 9.9 and 12.1 cases per 100 000 population over this time period.

Figure 1. Yersiniosis notifications by year, 2002-2006



There was some seasonality to the reporting of yersiniosis with more cases reported in the spring/summer months of October, November and January than other months over the five year period (Figure 1).

Figure 2. Average number of cases reported per month, 2002 to 2006



Over the five year period 2002 to 2006, the West Coast DHB had a yersiniosis notification rate significantly higher than the overall New Zealand rate every year. South Canterbury and Capital and Coast DHBs had a rate of yersiniosis significantly higher than the New Zealand rate in 2005 and 2006. Northland DHB consistently experienced a yersiniosis notification rate significantly less than the New Zealand rate.

Table 5. Number of cases and rates per 100,000 of yersiniosis notifications by DHB

| DHB | Year | | | | | | | | | |
|--------------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | Rate | No. | Rate | No. | Rate | No. | Rate | No. | Rate |
| Northland | 5 | 3.4* | 3 | 2.0* | 4 | 2.7* | 6 | 4.0* | 9 | 6.0* |
| Waitemata | 70 | 15.2 | 78 | 16.4* | 61 | 12.6 | 41 | 8.3 | 41 | 8.2* |
| Auckland | 67 | 16.7* | 56 | 13.5 | 57 | 13.5 | 47 | 11.0 | 45 | 10.5 |
| Counties Manukau | 44 | 10.9 | 23 | 5.5* | 34 | 8.0 | 31 | 7.1 | 34 | 7.7* |
| Waikato | 36 | 10.9 | 42 | 12.5 | 46 | 13.6 | 20 | 5.9* | 33 | 9.6 |
| Lakes | 12 | 12.0 | 10 | 9.9 | 7 | 6.9 | 9 | 8.9 | 20 | 19.7* |
| Bay of Plenty | 24 | 12.9 | 13 | 6.9 | 16 | 8.3 | 9 | 4.6* | 30 | 15.1 |
| Tairāwhiti | 3 | 6.6 | 15 | 33.2* | 5 | 11.1 | 4 | 8.9 | 4 | 9.0 |
| Taranaki | 3 | 2.8* | 5 | 4.7 | 6 | 5.7 | 5 | 4.8 | 7 | 6.7 |
| Hawke's Bay | 16 | 10.8 | 14 | 9.4 | 20 | 13.4 | 15 | 10.0 | 10 | 6.6 |
| Whanganui | 6 | 9.3 | 5 | 7.8 | 7 | 11.0 | 12 | 19.2* | 9 | 14.5 |
| MidCentral | 12 | 7.5 | 8 | 4.9* | 11 | 6.8 | 8 | 4.9* | 10 | 6.1* |
| Hutt | 10 | 7.3 | 17 | 12.3 | 23 | 16.6* | 6 | 4.3* | 8 | 5.8* |
| Capital and Coast | 38 | 14.6 | 34 | 12.8 | 36 | 13.3 | 48 | 17.5* | 69 | 24.7* |
| Wairarapa | 4 | 10.2 | 9 | 22.9* | 3 | 7.6 | 1 | 2.5 | 1 | 2.5 |
| Nelson Marlborough | 4 | 3.1* | 8 | 6.1 | 4 | 3.0* | 15 | 11.1 | 18 | 13.2 |
| West Coast | 20 | 64.9* | 13 | 42.3* | 14 | 45.8* | 16 | 52.4* | 9 | 29.5* |
| Canterbury | 65 | 14.5 | 58 | 12.7 | 40 | 8.6 | 69 | 14.6* | 89 | 18.6* |
| South Canterbury | 10 | 18.6 | 8 | 14.9 | 11 | 20.4* | 9 | 16.7 | 10 | 18.7 |
| Otago | 18 | 10.1 | 15 | 8.3 | 12 | 6.6 | 26 | 14.3 | 27 | 14.8 |
| Southland | 9 | 8.4 | 5 | 4.6 | 3 | 2.8* | 10 | 9.2 | 4 | 3.7* |
| New Zealand | 476 | 12.1 | 439 | 10.9 | 420 | 10.3 | 407 | 9.9 | 487 | 11.8 |

* significant difference between regional rate and annual NZ rate

The notification rate for children aged less than 1 year old and 1 to 4 years old was significantly higher than the overall New Zealand reported notification rate of yersiniosis in each of the five years (Table 6). Notification rates of yersiniosis in the 15-19 years age group were significantly lower than the overall New Zealand population each year.

Table 6. Number and rates per 100,000 of notifiable yersiniosis cases by age group

| Age group | Year | | | | | | | | | |
|-------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | Rate | No. | Rate | No. | Rate | No. | Rate | No. | Rate |
| <1 | 30 | 55.7* | 41 | 74.2* | 36 | 62.2* | 25 | 43.9* | 28 | 49.0* |
| 1 to 4 | 104 | 46.1* | 91 | 40.6* | 111 | 49.4* | 79 | 35.1* | 85 | 37.7* |
| 5 to 9 | 21 | 7.2* | 22 | 7.5 | 29 | 10.0 | 14 | 4.8* | 10 | 3.5* |
| 10 to 14 | 27 | 8.8 | 18 | 5.8* | 18 | 5.8* | 21 | 6.8 | 12 | 4.0* |
| 15 to 19 | 14 | 4.9* | 18 | 6.1* | 13 | 4.3* | 12 | 3.9* | 14 | 4.5* |
| 20 to 29 | 57 | 11.0 | 43 | 8.1 | 39 | 7.3* | 50 | 9.2 | 60 | 10.9 |
| 30 to 39 | 55 | 9.2 | 39 | 6.5* | 43 | 7.2* | 36 | 6.1* | 69 | 11.8 |
| 40 to 49 | 67 | 11.7 | 56 | 9.5 | 48 | 7.9 | 58 | 9.5 | 60 | 9.7 |
| 50 to 59 | 47 | 10.5 | 49 | 10.6 | 38 | 8.0 | 43 | 8.8 | 63 | 12.6 |
| 60 to 69 | 23 | 7.6* | 29 | 9.5 | 19 | 6.0* | 24 | 7.4 | 37 | 10.9 |
| 70+ | 28 | 8.4 | 31 | 9.1 | 17 | 4.9* | 35 | 9.9 | 46 | 12.8 |
| Unknown | 3 | | 2 | | 9 | | 10 | | 3 | |
| Total Cases | 476 | 12.1 | 439 | 10.9 | 420 | 10.3 | 407 | 9.9 | 487 | 11.8 |

* significant difference between age group and overall annual rate

3.7 Source of notifications

The majority of notifications over the period 2002 to 2006 were recorded in EpiSurv as originating from General Practitioners (GPs) although this percentage decreased from 87.2% in 2002 to 79.1% in 2006 (Table 7). All other reporting sources (hospital practitioners, laboratories and other sources) increased slightly over this time period.

Table 7. Number and percentage of notifiable yersiniosis cases by reporting source

| Reporting Source | Year | | | | | | | | | |
|-----------------------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| General practitioner | 415 | 87.2 | 372 | 84.7 | 346 | 82.4 | 315 | 77.4 | 385 | 79.1 |
| Hospital based practitioner | 25 | 5.3 | 30 | 6.8 | 32 | 7.6 | 33 | 8.1 | 48 | 9.9 |
| Laboratory | 29 | 6.1 | 33 | 7.5 | 32 | 7.6 | 47 | 11.5 | 41 | 8.4 |
| Other | 7 | 1.5 | 4 | 0.9 | 10 | 2.4 | 12 | 2.9 | 13 | 2.7 |
| Total | 476 | 100.0 | 439 | 100.0 | 420 | 100.0 | 407 | 100.0 | 487 | 100.0 |

3.8 Cases meeting yersiniosis case definition

Analysis of the case status data recorded in EpiSurv showed a very high percentage of cases were reported as meeting the criteria for a confirmed case of yersiniosis each year (Table 8).

Table 8. Number and percentage of notifiable yersiniosis cases by case status

| Case Status | Year | | | | | | | | | |
|-------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Confirmed | 472 | 99.2 | 429 | 97.7 | 412 | 98.1 | 402 | 98.8 | 476 | 97.7 |
| Probable | 1 | 0.2 | 0 | 0.0 | 1 | 0.2 | 1 | 0.2 | 1 | 0.2 |
| Unknown | 3 | 0.6 | 10 | 2.3 | 7 | 1.7 | 4 | 1.0 | 10 | 2.1 |
| Total | 476 | 100.0 | 439 | 100.0 | 420 | 100.0 | 407 | 100.0 | 487 | 100.0 |

Over the five year study period a high percentage of the notified cases were consistently recorded by PHS staff in EpiSurv as meeting the clinical criteria for a case of yersiniosis (Table 9).

Table 9. Number and percentage of notifiable confirmed yersiniosis cases reported meeting clinical criteria of case definition

| Fits clinical description | Year | | | | | | | | | |
|---------------------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Yes | 442 | 92.9 | 385 | 87.7 | 378 | 90.0 | 386 | 94.8 | 471 | 96.7 |
| No | 1 | 0.2 | 3 | 0.7 | 0 | 0.0 | 4 | 1.0 | 1 | 0.2 |
| Unknown | 33 | 6.9 | 51 | 11.6 | 42 | 10.0 | 17 | 4.2 | 15 | 3.1 |
| Total | 476 | 100.0 | 439 | 100.0 | 420 | 100.0 | 407 | 100.0 | 487 | 100.0 |

Two yersiniosis cases (one each from Tauranga and Rotorua) were de-notified (i.e. made not a case in EpiSurv). The reason for de-notification in both cases was a lack of clinical symptoms consistent with yersiniosis.

3.9 Enteric Reference Laboratory (ERL) *Yersinia* biotyping data

ERL is the only New Zealand laboratory that tests *Yersinia* isolates to determine the strain and biotype. The ERL dataset was therefore used as the definitive *Yersinia* strain and biotyping dataset for this project. Although only *Y. enterocolitica* and *Y. pseudotuberculosis* are notifiable, the strains for all isolates referred to ERL are shown and analysed in this section.

The number of isolates referred to ERL increased by 50% in the five year period from 107 in 2002 to 169 in 2006 (Table 10). The number of *Y. enterocolitica* isolates increased over this time period but the percentage of *Y. enterocolitica* isolates as a proportion of all isolates decreased over the five year period from 95% of isolates in 2002 to 77% in 2006. Over the same period the number of *Y. frederiksenii* isolates increased from 3% to 20% of all isolates.

Table 10. Number of isolates typed by ERL by strain of *Yersinia* as a percentage of all *Yersinia* isolates typed that year.

| <i>Yersinia</i> strain | Year | | | | | | | | | |
|----------------------------------|------|-----|------|-----|------|-----|------|-----|------|-----|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| <i>bercovieri</i> | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 |
| <i>enterocolitica</i> | 102 | 95 | 88 | 93 | 75 | 83 | 106 | 79 | 127 | 75 |
| <i>frederiksenii</i> | 3 | 3 | 3 | 3 | 9 | 10 | 21 | 16 | 33 | 20 |
| <i>intermedia</i> | 2 | 2 | 2 | 2 | 1 | 1 | 6 | 4 | 3 | 2 |
| <i>kristensenii</i> | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 1 | 1 |
| <i>pseudotuberculosis</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 1 |
| <i>rohdei</i> | 0 | 0 | 1 | 1 | 2 | 2 | 0 | 0 | 1 | 1 |
| unknown | 0 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | 1 | 1 |
| Negative for <i>Yersinia</i> spp | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 1 |
| Total | 107 | 100 | 95 | 100 | 90 | 100 | 135 | 100 | 169 | 100 |

The numbers and relative percentages of different *Y. enterocolitica* biotypes identified changed over the last five years (Table 11). In 2002 biotype 4 (42 cases) was the most common biotype identified followed by biotype 1A and 2. In 2006 biotype 1A (54 cases) was the most commonly reported type followed by biotype 4 (40 cases) and in equal numbers biotypes 2 and 3. Biotype 1A increased from 28 cases in 2002 to 54 cases in 2006. Biotypes 1B and 5 were rarely seen.


Table 11. Number and percentage of isolates typed by ERL by biotype of *Y. enterocolitica*

| <i>Y. enterocolitica</i> biotype | Year | | | | | | | | | |
|----------------------------------|------|----------------|------|-----|------|-----|------|-----|------|-----|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % ^a | No. | % | No. | % | No. | % | No. | % |
| 1A | 28 | 27 | 38 | 43 | 29 | 39 | 46 | 43 | 54 | 43 |
| 1B | 1 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 1 | 1 |
| 2 | 26 | 25 | 19 | 22 | 9 | 12 | 15 | 14 | 16 | 13 |
| 3 | 5 | 5 | 3 | 3 | 2 | 3 | 19 | 18 | 16 | 13 |
| 4 | 42 | 41 | 27 | 31 | 34 | 45 | 25 | 24 | 40 | 31 |
| 5 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 102 | 100 | 88 | 100 | 75 | 100 | 106 | 100 | 127 | 100 |

^a% - percentage of all *Y. enterocolitica* isolates sent to ERL in that year


Using the pathogenic groupings discussed earlier the *Y enterocolitica* biotype 1A cases represented a significant percentage (43%) of all reported cases typed in 2006 and the non pathogenic group of *Yersinia* cases represent an increasing percentage (3% in 2002 and 20% in 2006) of all typed isolates since mid 2004 (Figure 3). The number of cases that were pathogenic *Y enterocolitica* biotypes remained similar throughout the time period.

Figure 3. Graph of notifications by pathogenic group of *Yersinia* by year, 2002-2006



The pathogenic groupings were also used to examine variation in the biotypes by year and quarter. There was clustering of the different pathogenic groups in different quarters of the year (Figure 4).

Figure 4. Graph of *Yersinia* isolates by year, quarter and pathogenic grouping, 2002-2006



In order to carry out analysis by DHB and audit the notification data, *Yersinia* biotyping data was obtained for EpiSurv notifications by matching the records as described in the methods section. Table 12 shows the percentage of isolates matched to EpiSurv cases. The percentage of ERL *Yersinia* isolates matched to EpiSurv yersiniosis cases increased from 69% in 2002 to 85% in 2006.

Table 12. Number and percentage of isolates typed by ERL and matched to an EpiSurv case

| Isolates | 2002 | | 2003 | | Year 2004 | | 2005 | | 2006 | |
|--|------|----|------|----|--------------|----|------|----|------|----|
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Received by ERL for biotyping ^a | 107 | | 95 | | 90 | | 135 | | 169 | |
| Matched to an EpiSurv case | 72 | 67 | 74 | 78 | 67 | 74 | 108 | 80 | 143 | 85 |

^a Duplicate isolates for an individual case have been removed

More than 50% of isolates referred to ERL each year for notified diseases came from laboratories in the Canterbury and Otago DHBs (Table 13). Very few *Yersinia* isolates were sent to ERL from laboratories in other DHBs for typing although 2006 saw an increase for several DHBs (Waikato, Northland and Nelson Marlborough).

Table 13. Number of isolates typed by ERL and matched to an EpiSurv case as a percentage of all isolates typed that year, by DHB

| DHB | 2002 | | 2003 | | Year 2004 | | 2005 | | 2006 | |
|--------------------|------|----------------|------|----------------|--------------|----------------|------|----------------|------|----------------|
| | No. | % ^a | No. | % ^a | No. | % ^a | No. | % ^a | No. | % ^a |
| Northland | 2 | 3 | 0 | 0 | 0 | 0 | 5 | 5 | 9 | 6 |
| Waitemata | 1 | 1 | 1 | 1 | 3 | 4 | 5 | 5 | 1 | 1 |
| Auckland | 3 | 4 | 2 | 3 | 5 | 7 | 8 | 7 | 1 | 1 |
| Counties Manukau | 5 | 7 | 0 | 0 | 2 | 3 | 2 | 2 | 0 | 0 |
| Waikato | 2 | 3 | 6 | 8 | 6 | 9 | 2 | 2 | 13 | 9 |
| Lakes | 1 | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 1 | 1 |
| Bay of Plenty | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 2 |
| Tairāwhiti | 2 | 3 | 2 | 3 | 0 | 0 | 0 | 0 | 2 | 1 |
| Hawke's Bay | 4 | 6 | 3 | 4 | 10 | 15 | 0 | 0 | 4 | 3 |
| Taranaki | 2 | 3 | 2 | 3 | 1 | 1 | 0 | 0 | 2 | 1 |
| MidCentral | 0 | 0 | 2 | 3 | 1 | 1 | 2 | 2 | 3 | 2 |
| Whanganui | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 1 |
| Capital and Coast | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Hutt Valley | 0 | 0 | 0 | 0 | 2 | 3 | 1 | 1 | 0 | 0 |
| Wairarapa | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Nelson Marlborough | 1 | 1 | 3 | 4 | 1 | 1 | 3 | 3 | 12 | 8 |
| West Coast | 1 | 1 | 0 | 0 | 0 | 0 | 6 | 6 | 8 | 6 |
| Canterbury | 26 | 36 | 33 | 44 | 19 | 28 | 43 | 40 | 60 | 42 |
| South Canterbury | 1 | 1 | 2 | 3 | 2 | 3 | 2 | 2 | 0 | 0 |
| Otago | 18 | 25 | 15 | 20 | 13 | 19 | 25 | 23 | 23 | 16 |
| Southland | 2 | 3 | 2 | 3 | 1 | 1 | 4 | 4 | 0 | 0 |
| Total | 72 | 100 | 74 | 100 | 67 | 100 | 108 | 100 | 143 | 100 |

^a% - percentage of all isolates sent to ERL in that year

Most of the non pathogenic strains of *Yersinia* that were identified in isolates referred to ERL came from South Island DHBs (Table 14). Canterbury and Otago DHBs contributed 275 (59%) of the

Table 14. Number and percentage of *Yersinia* isolates typed by ERL and matched to an EpiSurv case that meet the current laboratory criteria for yersiniosis notification (*Y. enterocolitica* or *Y. pseudotuberculosis*) by DHB

| DHB | Year | | | | | | | | | | | | | | |
|--------------------|----------------|----------------------------|-----------|----------------|----------------------------|-----------|----------------|----------------------------|-----------|----------------|----------------------------|-----------|----------------|----------------------------|-----------|
| | 2002 | | | 2003 | | | 2004 | | | 2005 | | | 2006 | | |
| | Total isolates | Meet notification criteria | | Total isolates | Meet notification criteria | | Total isolates | Meet notification criteria | | Total isolates | Meet notification criteria | | Total isolates | Meet notification criteria | |
| No. | No. | % | No. | No. | % | No. | No. | % | No. | No. | % | No. | No. | % | |
| Northland | 2 | 2 | 100 | | | | | | 5 | 5 | 100 | 9 | 9 | 100 | |
| Waitemata | 1 | 1 | 100 | 1 | 1 | 100 | 3 | 3 | 100 | 5 | 5 | 100 | 1 | 1 | 100 |
| Auckland | 3 | 3 | 100 | 2 | 2 | 100 | 5 | 5 | 100 | 8 | 8 | 100 | 1 | 1 | 100 |
| Counties Manukau | 5 | 5 | 100 | | | | 2 | 2 | 100 | 2 | 2 | 100 | | | |
| Waikato | 2 | 2 | 100 | 6 | 6 | 100 | 6 | 5 | 83 | 2 | 2 | 100 | 13 | 13 | 100 |
| Lakes | 1 | | | | | | 1 | 1 | 100 | | | | 1 | 1 | 100 |
| Bay of Plenty | | | | | | | | | | | | | 3 | 3 | 100 |
| Tairāwhiti | 2 | 2 | 100 | 2 | 2 | 100 | | | | | | | 2 | 2 | 100 |
| Hawke's Bay | 4 | 4 | 100 | 3 | 3 | 100 | 10 | 10 | 100 | | | | 4 | 3 | 75 |
| Taranaki | 2 | 2 | 100 | 2 | 2 | 100 | 1 | 1 | 100 | | | | 2 | 2 | 100 |
| MidCentral | | | | 2 | 2 | 100 | 1 | 1 | 100 | 2 | 2 | 100 | 3 | 3 | 100 |
| Whanganui | 1 | 1 | 100 | 1 | 0 | - | | | | | | | 1 | 0 | - |
| Capital and Coast | | | | | | | | | | | | | | | |
| Hutt Valley | | | | | | | 2 | 2 | 100 | 1 | 1 | 100 | | | |
| Wairarapa | | | | | | | | | | | | | | | |
| Nelson Marlborough | 1 | 1 | 100 | 3 | 3 | 100 | 1 | 1 | 100 | 3 | 3 | 100 | 12 | 11 | 92 |
| West Coast | 1 | 1 | 100 | | | | | | | 6 | 6 | 100 | 8 | 5 | 63 |
| Canterbury | 26 | 23 | 88 | 33 | 33 | 100 | 19 | 12 | 63 | 43 | 32 | 74 | 60 | 38 | 63 |
| South Canterbury | 1 | 1 | 100 | 2 | 2 | 100 | 2 | 2 | 100 | 2 | 2 | 100 | | | |
| Otago | 18 | 18 | 100 | 15 | 13 | 87 | 13 | 9 | 69 | 25 | 14 | 56 | 23 | 17 | 74 |
| Southland | 2 | 2 | 100 | 2 | 2 | 100 | 1 | 0 | - | 4 | 2 | 50 | | | |
| Total | 72 | 68 | 94 | 74 | 71 | 96 | 67 | 54 | 83 | 108 | 84 | 78 | 143 | 109 | 76 |

464 isolates over the 5 year period but accounted for 68 (87%) of the 78 non pathogenic strains of *Yersinia* isolated.

Analysis of the combined yersiniosis notification and biotyping data for the five year period showed an increasing number of isolates being referred to ERL. However the percentage of these isolates that met the notification criteria and the percentage of the isolates that were found to be a pathogenic strain both decreased over the five year period. (Table 15).

Table 15. Summary of notifications and laboratory biotyping results, 2002 to 2006

| Year | Notifications | Samples sent to ERL ^a | | Meets notification criteria ^b | | Pathogenic strain | |
|------|---------------|----------------------------------|------|--|------|-------------------|------|
| | No. | No | % | No. | % | No | % |
| 2002 | 476 | 72 | 15.1 | 68 | 94.4 | 53 | 73.6 |
| 2003 | 439 | 74 | 16.9 | 71 | 95.9 | 41 | 55.4 |
| 2004 | 420 | 67 | 16.0 | 54 | 80.6 | 34 | 50.7 |
| 2005 | 407 | 108 | 26.5 | 84 | 77.8 | 48 | 44.4 |
| 2006 | 487 | 143 | 29.4 | 109 | 76.2 | 63 | 44.1 |

^aThis includes only samples that were matched to a disease notification

^b Includes only *Y. enterocolitica* or *Y. pseudotuberculosis* isolates

Further analysis of this trend for DHBs that submitted more than five samples to ERL in the five year period revealed low percentages of pathogenic strains of *Yersinia* were being identified for some DHBs e.g. Canterbury (33.7%), West Coast (26.7%), Southland (22.2%) (Table 16).

An audit of the *Yersinia* biotyping information entered into EpiSurv against the biotyping data that was available in the ERL database showed that a large percentage of the potentially available *Yersinia* biotyping information had not been entered into EpiSurv (Table 17). The records with *Yersinia* strain information entered increased from 14% (11/78) of cases in 2002 to 26.2% (38/145) of cases in 2006.

Table 16. Summary of notifications and laboratory biotyping results by DHB for the period 2002 to 2006

| DHB | Notifications | Samples sent to ERL ^a | | Meets notification criteria ^b | | Pathogenic strain | |
|-------------------|---------------|----------------------------------|------|--|-------|-------------------|-------|
| | No. | No | % | No. | % | No | % |
| Northland | 27 | 16 | 59.3 | 16 | 100.0 | 13 | 81.3 |
| Waitemata | 291 | 11 | 3.8 | 11 | 100.0 | 8 | 72.7 |
| Auckland | 274 | 19 | 6.9 | 19 | 100.0 | 16 | 84.2 |
| Counties Manukau | 164 | 9 | 5.5 | 9 | 100.0 | 9 | 100.0 |
| Waikato | 187 | 29 | 15.5 | 28 | 96.6 | 25 | 86.2 |
| Lakes | 57 | 3 | 5.3 | 2 | 66.7 | 1 | 33.3 |
| Bay of Plenty | 91 | 3 | 3.3 | 3 | 100.0 | 2 | 66.7 |
| Tairāwhiti | 31 | 6 | 19.4 | 6 | 100.0 | 4 | 66.7 |
| Hawke's Bay | 75 | 21 | 28.0 | 20 | 95.2 | 17 | 81.0 |
| Taranaki | 27 | 7 | 25.9 | 7 | 100.0 | 4 | 57.1 |
| MidCentral | 59 | 8 | 13.6 | 8 | 100.0 | 4 | 50.0 |
| Whanganui | 29 | 3 | 10.3 | 1 | 33.3 | 1 | 33.3 |
| Capital and Coast | 215 | 0 | 0.0 | 0 | - | 0 | - |
| Hutt Valley | 64 | 3 | 4.7 | 3 | 100.0 | 3 | 100.0 |
| Wairarapa | 18 | 0 | 0.0 | 0 | - | 0 | - |
| Nelson | | | | | | | |
| Marlborough | 49 | 20 | 40.8 | 19 | 95.0 | 14 | 70.0 |
| West Coast | 72 | 15 | 20.8 | 12 | 80.0 | 4 | 26.7 |
| Canterbury | 322 | 181 | 56.2 | 138 | 76.2 | 61 | 33.7 |
| South Canterbury | 48 | 7 | 14.6 | 7 | 100.0 | 3 | 42.9 |
| Otago | 105 | 94 | 89.5 | 71 | 75.5 | 48 | 51.1 |
| Southland | 24 | 9 | 37.5 | 6 | 66.7 | 2 | 22.2 |
| Total | 2229 | 464 | 20.8 | 386 | 83.2 | 239 | 51.5 |

^a Includes only samples that were matched to a disease notification

^b Includes only *Y. enterocolitica* or *Y. pseudotuberculosis* isolates

Table 17. Number of yersiniosis notifications reported on EpiSurv, *Yersinia* isolates typed by the ERL and EpiSurv records updated with *Yersinia* strain by DHB and year, 2002 to 2006.

| DHB | 2002 | | | 2003 | | | Year 2004 | | | 2005 | | | 2006 | | |
|--------------------|------------------|------------------|--------------------|------------|-----------|---------------|--------------|-----------|---------------|------------|------------|---------------|------------|------------|---------------|
| | Epi ^a | ERL ^b | Updtd ^c | Epi | ERL | Updtd | Epi | ERL | Updtd | Epi | ERL | Updtd | Epi | ERL | Updtd |
| Northland | 5 | 2 | | 3 | | | 4 | | | 6 | 5 | | 9 | 9 | 1(0) |
| Waitemata | 70 | 1 | | 78 | 1 | | 61 | 3 | | 41 | 5 | | 41 | 1 | |
| Auckland | 67 | 3 | | 58 | 2 | | 57 | 5 | | 47 | 8 | | 45 | 1 | |
| Counties Manukau | 44 | 5 | | 21 | | | 34 | 2 | | 31 | 2 | | 34 | | |
| Waikato | 38 | 2 | | 43 | 6 | | 47 | 6 | | 25 | 2 | | 34 | 13 | |
| Lakes | 12 | 1 | | 10 | | | 7 | 1 | | 9 | | | 19 | 1 | |
| Bay of Plenty | 23 | | | 13 | | | 16 | | | 9 | | | 30 | 3 | |
| Tairāwhiti | 3 | 2 | 2(1) | 15 | 2 | 1(0) | 5 | | | 4 | | | 4 | 2 | |
| Hawke's Bay | 16 | 4 | | 14 | 3 | | 20 | 10 | | 15 | | | 10 | 4 | |
| Taranaki | 3 | 2 | | 5 | 2 | | 6 | 1 | | 5 | | | 8 | 2 | |
| MidCentral | 14 | | | 10 | 2 | 1(0) | 11 | 1 | | 10 | 2 | | 14 | 3 | |
| Whanganui | 4 | 1 | | 4 | 1 | 1(1) | 6 | | | 7 | | | 8 | 1 | |
| Capital and Coast | 36 | | | 32 | | | 36 | | | 46 | | | 65 | | |
| Hutt Valley | 10 | | | 17 | | | 23 | 2 | | 6 | 1 | | 8 | | |
| Wairarapa | 4 | | | 9 | | | 3 | | | 1 | | | 1 | | |
| Nelson Marlborough | 4 | 1 | | 8 | 3 | | 4 | 1 | | 15 | 3 | | 18 | 12 | 2(2) |
| West Coast | 20 | 1 | | 13 | | | 14 | | | 16 | 6 | | 9 | 8 | |
| Canterbury | 66 | 26 | 9(5) | 58 | 33 | 15(12) | 40 | 19 | 11(10) | 69 | 43 | 23(17) | 89 | 60 | 34(33) |
| South Canterbury | 10 | 1 | | 8 | 2 | | 11 | 2 | 1(0) | 9 | 2 | | 10 | | |
| Otago | 21 | 18 | | 15 | 15 | | 14 | 13 | | 28 | 25 | 4(4) | 27 | 23 | 1(1) |
| Southland | 6 | 2 | | 5 | 2 | | 1 | 1 | | 8 | 4 | | 4 | | |
| Total | 476 | 72 | 11(6) | 439 | 74 | 18(13) | 420 | 67 | 12(10) | 407 | 108 | 27(21) | 487 | 143 | 38(36) |

^aEpi = number of notified cases reported on EpiSurv

^bERL = number of EpiSurv cases that could be matched to a *Yersinia* isolate typed by ERL

^cUpdtd = number of EpiSurv cases that have been updated with the strain (and biotype) of *Yersinia* as determined by ERL.

3.10 Hospitalisations

Two sources of hospitalisation data, EpiSurv notifications and NZHIS morbidity data were used to analyse trends in yersiniosis over the past five years.

The percentage of EpiSurv cases that were as hospitalised on EpiSurv increased from 6.3% in 2002 to 11.5% in 2006 (Table 18).

Table 18. Number and percentage of notified yersiniosis cases admitted to a hospital reported on EpiSurv

| Hospitalised | Year | | | | | | | | | |
|--------------|------|-------|------|-------|------|-------|------|-------|------|-------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Yes | 30 | 6.3 | 30 | 6.8 | 25 | 6.0 | 41 | 10.1 | 56 | 11.5 |
| No | 305 | 64.1 | 234 | 53.3 | 215 | 51.2 | 212 | 52.1 | 293 | 60.2 |
| Unknown | 141 | 29.6 | 175 | 39.9 | 180 | 42.9 | 154 | 37.8 | 138 | 28.3 |
| Total | 476 | 100.0 | 439 | 100.0 | 420 | 100.0 | 407 | 100.0 | 487 | 100.0 |

The number of cases reported to NZHIS with ICD-10 code A046 (yersiniosis) for both primary and other relevant diagnosis increased from 26 in 2002 to 50 in 2006 (Table 19).

Table 19. Number of hospitalised yersiniosis cases (ICD10 A046) by diagnosis status, as reported by NZHIS

| Type of diagnostic code | Year | | | | |
|--------------------------|------|------|------|------|------|
| | 2002 | 2003 | 2004 | 2005 | 2006 |
| Principal diagnosis | 14 | 7 | 17 | 12 | 27 |
| Other relevant diagnosis | 12 | 10 | 13 | 15 | 23 |
| Total | 26 | 17 | 30 | 27 | 50 |

The NZHIS hospitalisation records for yersiniosis were matched to the EpiSurv and ESRLab records as described in the methods section. Table 20 shows the number of records that were matched from these sources.

Table 20. Number and percentage of hospitalised yersiniosis reported by NZHIS that could be matched to cases on EpiSurv and isolates typed by ERL

| NZHIS records | Year | | | | | | | | | |
|--------------------|------|----|------|-----|------|----|------|----|------|----|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | No. | % | No. | % | No. | % | No. | % | No. | % |
| Matched to EpiSurv | 17 | 65 | 17 | 100 | 20 | 67 | 19 | 70 | 38 | 76 |
| Matched to ERL | 6 | 23 | 10 | 59 | 15 | 50 | 12 | 44 | 26 | 52 |
| NZHIS Total | 26 | - | 17 | - | 30 | - | 27 | - | 50 | - |

The ERL *Yersinia* strain and biotype data that was derived for the NZHIS hospitalised cases is shown in Table 21. The numbers were very small but a trend of an increasing number of cases of *Y. enterocolitica* 1A and *Y. frederiksenii* can be seen in recent years.

Table 21. *Yersinia* strain and biotype (from ERL) for NZHIS reported yersiniosis cases by diagnostic status

| <i>Yersinia</i> strain and biotype | Year | | | | | | | | | |
|------------------------------------|---------------------|--------------------------|---------------------|--------------------------|---------------------|--------------------------|---------------------|--------------------------|---------------------|--------------------------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | Principal diagnosis | Other relevant diagnosis | Principal diagnosis | Other relevant diagnosis | Principal diagnosis | Other relevant diagnosis | Principal diagnosis | Other relevant diagnosis | Principal diagnosis | Other relevant diagnosis |
| <i>enterocolitica</i> 1A | 1 | 1 | | 6 | 4 | 4 | 2 | 7 | 6 | 7 |
| <i>enterocolitica</i> 2 | 1 | | | | 1 | 1 | 1 | | 1 | |
| <i>enterocolitica</i> 3 | | 1 | | | | | | | | |
| <i>enterocolitica</i> 4 | 1 | | 3 | 1 | 2 | 1 | | | 2 | 1 |
| <i>frederiksenii</i> | | | | | | 1 | | 2 | 3 | 4 |
| <i>intermedia</i> | | 1 | | | | | | | | 1 |
| <i>rohdei</i> | | | | | 1 | | | | | |
| unknown | | | | | | | | | | 1 |
| Total | 3 | 3 | 3 | 7 | 8 | 7 | 3 | 9 | 12 | 14 |

Table 22 shows that with the exception of the 2004 reporting year more hospitalisations were reported on EpiSurv than the total number of cases hospitalised with yersiniosis as a primary or other level diagnostic code reported through NZHIS. Approximately half of the EpiSurv notifications matched an NZHIS yersiniosis record with yersiniosis as a primary diagnosis.

Over the past five years the number of hospitalisations reported in EpiSurv for each DHB was generally less than 5 with the exception of the Canterbury DHB which in 2006 accounted for 32 of the 56 hospitalisations (57%) reported for all of New Zealand.

Table 22. Number of hospitalised yersiniosis cases reported on EpiSurv compared to NZHIS reported cases.

| DHB | Year | | | | | | | | | |
|--------------------|------------------|----------------------|-----------|---------------|-----------|----------------|-----------|----------------|-----------|----------------|
| | 2002 | | 2003 | | 2004 | | 2005 | | 2006 | |
| | Epi ^a | HIS(1°) ^b | Epi | HIS (1°) | Epi | HIS (1°) | Epi | HIS (1°) | Epi | HIS (1°) |
| Northland | | | | | | 2 (1) | | 1 (0) | 3 | 4 (4) |
| Waitemata | 1 | 3 (1) | 1 | | 2 | 3 (2) | | | 1 | 2 (1) |
| Auckland | 1 | 3 (2) | 1 | 3 (3) | 1 | 2 (1) | 1 | 2 (2) | 2 | 3 (3) |
| Counties Manukau | 2 | 4 (2) | 2 | 2 (1) | | 4 (1) | 1 | 3 (1) | | 5 (4) |
| Waikato | | | 6 | 2 (2) | 3 | 1 (0) | 5 | | 1 | 1 (0) |
| Lakes | | 1 (1) | | | 1 | | | | 2 | 1 (1) |
| Bay of Plenty | 2 | 3 (3) | 2 | | 1 | 1 (1) | 2 | 1 (1) | 3 | 4 (1) |
| Tairāwhiti | 3 | | 1 | 1 (1) | 1 | 1 (0) | | | | |
| Hawke's Bay | | | 2 | | 2 | 2 (2) | 1 | | | 1 (0) |
| Taranaki | | 1 (0) | | | | | | 3 (1) | | |
| MidCentral | 1 | 2 (1) | 2 | 1 (0) | 1 | 1 (1) | 1 | 1 (1) | 1 | 2 (1) |
| Whanganui | | | | | | 1 (1) | 2 | 1 (1) | | |
| Capital and Coast | 5 | | | | 2 | | 7 | 4 (2) | 3 | 2 (0) |
| Hutt Valley | | | | | 1 | 1 (0) | | | 1 | 2 (2) |
| Wairarapa | | | | | | | 1 | | | |
| Nelson Marlborough | | 1 (1) | 1 | | 1 | 1 (1) | | | 3 | 2 (0) |
| West Coast | 4 | 3 (2) | 3 | 1 (0) | | 1 (1) | 1 | 1 (0) | 1 | 3 (0) |
| Canterbury | 8 | 5 (1) | 5 | 5 (0) | 6 | 5 (1) | 18 | 9 (3) | 32 | 14 (8) |
| South Canterbury | | | 1 | 1 (0) | | 1 (1) | | 1 (0) | | |
| Otago | 1 | | 1 | 1 (0) | 2 | 2 (2) | 1 | | 3 | 4 (2) |
| Southland | 2 | | 2 | | 1 | 1 (0) | | | | |
| Total | 30 | 26 (14) | 30 | 17 (7) | 25 | 30 (16) | 41 | 27 (12) | 56 | 50 (27) |

^aEpi - number of hospitalised yersiniosis cases reported on EpiSurv

^bHIS (1°) - total number of hospitalised yersiniosis cases (yersiniosis given as the principal diagnosis) reported by the NZHIS

3.11 *Yersinia* testing methodologies used by clinical laboratories in New Zealand

The Acute Gastrointestinal Illness (AGI) Study: Laboratory Survey of 46 community, hospital and public health laboratories across New Zealand, undertaken in mid-2006, investigated the criteria and laboratory methodologies used to test for various enteric pathogens including *Yersinia*. [6]

This survey, which had a 76% response rate, found that it was generally standard practice to include testing for *Yersinia* when undertaking an “enteric screen” on faecal specimens (32/34 respondents).

Approximately half of laboratories reported that they would refer on positive *Yersinia* isolates to the Enteric Reference Laboratory.

3.12 Medical Officers of Health Survey

Responses to the survey were received from every Public Health office with a Medical Officer of Health..

3.12.1 Source of notification

All PHSs reported receiving yersiniosis notifications from GPs. Fifteen reported receiving notifications from hospital clinicians and eight received notifications direct from laboratories. For those receiving direct laboratory notifications five (Wanganui, Canterbury, Tairāwhiti, Hawke's Bay and Blenheim) had been receiving results for more than five years and three (MidCentral, Northland and Taranaki) started receiving results in the last five years.

Four PHSs reported having carried out an audit of laboratory data against notification data to assess whether all cases are notified. The proportion of all yersiniosis cases notified varied from 70% in Auckland to 90% in Tauranga and 100% in Nelson and Otago.

All PHSs except Northland and Canterbury reported entering all yersiniosis notifications onto EpiSurv. Northland reported entering only those cases that met the case definition. A respondent from Canterbury PHS reported that "as per the Public Health Surveillance Manual only *Y. enterocolitica* and *Y. pseudotuberculosis* are entered as cases. However if we do not have the type of *Yersinia* at the time of notification they are all entered. When the typing comes through this is updated on EpiSurv and made "not a case" if they are any type other than those mentioned above."

3.12.2 Investigation of cases

All PHSs except Auckland investigate all of the yersiniosis cases that are notified. Auckland will investigate a geographic cluster of cases and when two or more cases are potentially linked to a common source.

Investigation of yersiniosis cases is undertaken by Health Protection Officers (or PHS Designated Officers) in 12 of the 17 PHSs. Environmental Health Officers are involved in the investigation of yersiniosis notification in five PHSs and EpiSurv Co-ordinators were responsible for investigation in two PHSs.

Nine PHSs use a specific questionnaire and seven of these have developed their own questionnaire.

Fourteen PHSs make a phone call to the case as part of the investigation. Six PHSs post out questionnaires with return rates ranging from 57 to 95%. Two PHSs administer the questionnaire in person.

Ten PHSs confirmed that the notified cases had met the yersiniosis clinical description with the case or notifying practitioner. The policy for two PHSs was to

confirm that the case met the clinical criteria during the case investigation and the remaining six PHSs reported having no policy.

3.12.3 Local trends in yersiniosis notifications

Twelve PHSs had not noticed any trends in yersiniosis notifications over the past 1-2 years. Four PHSs (Tauranga, Capital and Coast, West Coast and Canterbury) had noticed recent increases and one (Auckland) had seen occasional clusters.

3.12.4 Yersinia strain and biotyping information

Only one PHS (MidCentral) reported always receiving species information. Ten PHSs reported sometimes receiving information on the species of *Yersinia* and four PHSs (Auckland, Hamilton, Nelson and Hawke's Bay) reported never receiving species information. Two PHSs did not know or did not answer the question.

Of the PHSs receiving information on the species of *Yersinia* (10 PHSs), this information came from the local laboratory (6/10 Tauranga, Tairāwhiti, MidCentral, Wanganui, Canterbury, Otago), the GP (4/10 Tauranga, West Coast, Canterbury, South Canterbury) and ERL (4/10, Tairāwhiti, Capital and Coast, West Coast, Canterbury).

For four PHSs (Tauranga, West Coast, Canterbury and Blenheim) it was their policy to enter all *Yersinia* species results received onto EpiSurv. The policy for the remainder varied. Some only entered the "unusual species i.e. the non *Y. enterocolitica* species".

3.12.5 De-notification of yersiniosis cases

De-notification of a yersiniosis case (i.e. changing the status in EpiSurv to "not a case") was never considered by eight PHSs.

The remaining considered de-notification if subsequently advised by the laboratory or GP that the case did not meet the clinical symptoms or a negative *Yersinia* laboratory result was reported (6 PHSs) or the case did not meet the case definition as per the *Communicable Disease Control* manual (3).

3.12.6 Other Comments

One respondent asked for "more information as to what the pathogenic and non pathogenic strains are as apparently *Y. enterocolitica* biotype 1A is also non pathogenic which would make up the bulk of our notifications. The typing is received from the hospital lab after they receive it from the ESR ERL."

4 DISCUSSION

The number of yersiniosis notifications reported increased from 2002 to 2006. Most of this increase appeared to be from South Island DHBs and particularly Canterbury DHB. Despite the low population of the West Coast DHB it consistently reported higher population rates of yersiniosis notifications than other DHBs and sufficient numbers were reported for stability of rates to be an unlikely explanation for the higher rates.

Only a small percentage of all notifications were typed by ERL. The practice of referring isolates for biotyping varies throughout the country with clinical laboratories in some DHBs (e.g. Capital and Coast) referring no isolates to ERL in the five year period and laboratories in other DHBs (e.g. Canterbury) referring a high percentage of isolates.

The percentage of notified cases with isolates referred to ERL increased in the last two years with most of this increase found to be due to non pathogenic subtypes e.g. *Y. frederiksenii*.

The case definition for yersiniosis includes both isolation of *Y. enterocolitica* or *Y. pseudotuberculosis* from blood or faeces and detection of circulating antigen by ELISA or agglutination test. In the AGI study no laboratories were found to be routinely carrying out the latter test and this was never noted in EpiSurv as the test undertaken for the notification in time period analysed.

Given that isolation is the primary means of identification for *Yersinia* and only ERL carries out strain and biotyping of *Yersinia* to determine whether the cases are *Y. enterocolitica* or *Y. pseudotuberculosis*. For 70% of notifications it was not possible to say whether the case met the yersiniosis case definition. Further study of a DHB with high numbers of cases typed (e.g. Canterbury) may further inform this question but it is difficult to generalise this information to other DHBs where no typing had been undertaken.

There was a large amount of variation in the percentage of pathogenic strains of *Yersinia* being isolated in the samples submitted by clinical laboratories from different DHBs. Of most concern is the increasing numbers of samples being submitted to ERL and the increasing percentage that are either not pathogenic or *Yersinia enterocolitica* biotype 1A. It would be useful to examine in greater detail the methods being used by different laboratories to isolate *Yersinia*. The Acute Gastrointestinal Illness (AGI) Study: Laboratory Survey provided insufficient detail to address this issue.

Although some PHSs indicated they would de-notify yersiniosis cases where the case definition was not met, this did not appear to happen very often.

Over the past three years ESR has been matching many of the ERL isolates to EpiSurv records. This information has been used to identify non notified cases of disease where ESR carries out typing of all isolates (e.g. salmonellosis and meningococcal disease) and the PHS is provided with information to follow up the disease cases not notified. With diseases such as salmonellosis, laboratories refer almost all samples to ERL for typing. ESR matches the ESRLab data with EpiSurv

notifications and requests PHSs to de-notify cases where the laboratory criteria have not been met. This is more difficult for yersiniosis given the small percentage of isolates referred to ERL and therefore the many yersiniosis notifications for which the strain is unknown.

When the *Yersinia* isolates have been biotyped by ERL, the results are sent back to the requesting clinical laboratory but many of the results do not appear to have been received by the PHSs and updated on EpiSurv. Many PHSs in the MOsH survey stated that if they received laboratory test results these would be added to the appropriate notification records on EpiSurv. The new requirement (implemented from 18 December 2007) under the Health Act for laboratories to directly report notifiable diseases to Medical Officers of Health will allow ERL to send biotyping data for yersiniosis cases directly to PHSs.

In addition to informing PHSs of non notified cases of disease, since the implementation of the web based EpiSurv in April 2007 the ESR held organism typing information for selected diseases (e.g. *Salmonella*, *Shigella*) is now updated by ESR directly into EpiSurv. The organism typing information for *Yersinia* in EpiSurv could be added to this process.

Biotyping data was rarely used by PHS staff even in PHSs where typing data was available. There was uncertainty about how to use the biotyping data to inform investigations and in particular which strains and biotypes of *Yersinia* are pathogenic or non pathogenic.

Given the issue of pathogenicity of different strains of *Yersinia* the biotyping data is an essential tool for the investigation of clusters of yersiniosis cases. Monitoring of the biotyping data at a national and regional level would ensure that clusters are identified. Better use of the biotyping data would also assist PHSs to prioritise the investigation of yersiniosis cases and look for common sources amongst the same pathogenic biotypes.

The number of records in some of the matched datasets (e.g. hospitalisations) was very small and therefore suffered from the usual issues of instability with small numbers. However there was a clear recent increasing trend in the notifications for hospitalised cases from the Canterbury DHB area that would benefit from further investigation.

The larger number of reported hospitalisations among EpiSurv yersiniosis cases than reported by NZHIS suggests that a clinical diagnosis of yersiniosis may not always be present. Further investigation by the PHS staff is needed to understand this situation better.

5 RECOMMENDATIONS

It is recommended that

1) NZFSA communicates the findings of this report to MoH and other interested parties e.g. PHSs to consider the development of the following enhancements for *Yersinia* surveillance

a) Development of guidelines for the use of *Yersinia* biotyping data in yersiniosis investigations by PHS staff

b) Development of a consistent policy for clinical laboratories to refer *Yersinia* isolates to ESR for *Yersinia* biotyping.

c) Development of a policy for the recording and updating of the case status (e.g. confirmed, not a case) in EpiSurv on receipt of *Yersinia* laboratory test results

d) Development of aberration reporting to detect time and space clusters of *Y. enterocolitica* biotypes.

2) ESR consider adding yersiniosis to the list of disease records that are matched between ESRLab and EpiSurv and update strain and biotyping data for yersiniosis notifications directly into EpiSurv.

3) Canterbury DHB further investigates the recent increase in the Canterbury DHB and the high rates in the South Canterbury and West Coast DHBs using available biotyping data.

4) ESR reviews the methods for typing of *Yersinia* isolates and considers options for improving the turnaround time for biotyping of results.

6 REFERENCES

1. Cressey, P. and R. Lake, *Risk Ranking: Estimates of the Burden of Foodborne Disease for New Zealand*. June 2007, ESR Client Report FW0724: Christchurch, NZ.
2. Rocourt, J., et al., *The present state of foodborne disease in OECD countries*. 2003, World Health Organization: Geneva.
3. *Communicable Disease Control Manual*. June 1988, Public Health Group Ministry of Health.
4. Robins-Browne, R.M., *Yersinia enterocolitica*, in *Food microbiology : fundamentals and frontiers, 3rd Ed.*, M.P. Doyle and L.R. Beuchat, Editors. 2007, ASM Press: Washington, D.C. p. 293-322.
5. Tennant, S.M., T.H. Grant, and R.M. Robins-Browne, *Pathogenicity of Yersinia enterocolitica biotype 1A*. FEMS Immunology and Medical Microbiology, 2003. **38**: p. 127-137.
6. *Acute Gastrointestinal Illness (AGI) Study: Laboratory Survey*, in *Client report FW0685, prepared for the New Zealand Food Safety Authority*. 2007, Institute of Environmental Science and Research Limited: Christchurch.

APPENDIX A – MEDICAL OFFICERS OF HEALTH YERSINIOSIS SURVEY RESPONDENTS

| Public Health Office | Respondent |
|---------------------------------------|---------------------|
| Northland | Jonathan Jarman |
| Auckland, Waitemata, Counties Manukau | Craig Thornley |
| Waikato | Maureen O'Halloran |
| Bay of Plenty | Lynnette Borissenko |
| Tairāwhiti | Alan Hall |
| Taranaki | Viv O'Leary |
| Hawke's Bay | Paul Buckley |
| MidCentral | Peter Wood |
| Wanganui | Margaret Tunbridge |
| Capital & Coast, Hutt Valley | Quentin Ruscoe |
| Nelson | Leanne Punt |
| Marlborough | Kirsten Todd |
| Canterbury | Debbie Smith |
| West Coast | Cheryl Brunton |
| South Canterbury | Daniel Williams |
| Otago | Kaylene Newell |
| Southland | Chris McCall |

APPENDIX B – YERSINIOSIS QUESTIONNAIRE

Public Health Service Yersiniosis Notifications Questionnaire

Yersiniosis is the third commonest potential food-borne disease notified in New Zealand. New Zealand's notification rate for yersiniosis (11.8 cases per 100,000 in 2006) is high compared to many other countries.

NZFSA has asked ESR to review the yersiniosis data for the past five years. A review of yersiniosis notifications reported on EpiSurv, laboratory data held by ESR's Enteric Reference Laboratory and yersiniosis hospitalisation data from NZHIS is currently being carried out.

To assist in validating and interpreting the EpiSurv data we would appreciate if **one person from each Public Health Service**, or Public Health Office if each office has different reporting procedures, completes this questionnaire and return to Jonathan Williman jonathan.williman@esr.cri.nz at ESR by the 5pm, Wednesday 14th November 2007.

Public Health Service / Office: _____

Name of person completing questionnaire: _____

Please check appropriate answer. Text boxes will expand as you type.

1. In your PHS where do yersiniosis notifications come from? (*tick all that apply*)

- GP
- Hospital clinician
- Directly from local laboratory (e.g. hospital or community laboratory)

If yes to direct laboratory notification please answer these questions

a) When did direct laboratory notification start (approximately)?

b) Which isolations of *Yersinia* are you notified of?

- All positive isolations
- Only positive isolations where case is known to have clinical illness
- Other _____
- Other source (*please specify*) _____

2. Has your PHS ever audited laboratory data against notification data to assess whether all cases (i.e. positive laboratory result and clinical illness present) are notified?

- Yes No Don't know

If yes, what proportion of all yersiniosis cases are notified? _____%

3. Are all notifications of yersiniosis that your PHS receives entered onto EpiSurv?

- Yes
 No *please give details of types of notifications not entered* _____

4. Which yersiniosis cases do you investigate¹? All Some None Don't know

If you answered None or Don't know please go to Question 5

a) Please outline the criteria used by your PHS for deciding which cases will be investigated

b) What percentage of cases do you investigate? _____

c) Who undertakes the yersiniosis investigations?

- HPOs EHOs Other *please specify* _____

d) Is a specific yersiniosis questionnaire used?

- Yes No Don't know

If yes, did your PHS develop this or was it acquired from somewhere else?

- Developed by PHS Acquired from _____

e) How is the questionnaire administered?

- Phone call to case
 Posted out

If posted out please answer these questions

i) Is the postal questionnaire followed up if not returned

- Yes No Don't know

ii) What percentage of postal questionnaires are returned?
_____%

- Other *please specify* _____

¹ Contact with case through questionnaire, phone call or visit

f) What is it the policy at your PHS for confirming that notified cases meet the yersiniosis clinical description?

(e.g. no policy, ask notifying practitioner, confirm with case)

5. Have you noticed any particular local trends in yersiniosis notifications over the last 1-2 years (e.g. increase or decrease in incidence, demographics of cases, risk factors, sources of notification)?

Yes please give details _____

No

6. Does your PHS receive information regarding the particular species of *Yersinia*?

Yes, always
Don't know

Yes, sometimes

No, never

If yes

a) Where does the information regarding the *Yersinia* species come from?

GP

Local laboratory

ESR Enteric Reference Laboratory

ESR Population and Environmental Health Group

b) Is it the policy at your PHS to enter all *Yersinia* species results received onto EpiSurv (i.e. into the Additional Laboratory Results field)?

Yes

No

Don't know

7. Under what circumstances would you consider de-notification of a yersiniosis case (i.e. changing the status in EpiSurv to "not a case")?

8. Do you have any other comments regarding yersiniosis notifications?
